


MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
Effective Date: 10/1/2022	Medical Policy Number: 101
 10/1/2022	Medical Policy Committee Approved Date: 9/05; 5/07; 11/07; 11/08; 12/10; 10/12; 9/13; 10/14; 7/15; 9/15; 6/16; 8/16; 10/17; 12/18; 12/19; 5/2020; 07/2021; 7/2022; 9/2022
Medical Officer	Date

See Policy CPT CODE section below for any prior authorization requirements

SCOPE:

Providence Health Plan, Providence Health Assurance, Providence Plan Partners, and Ayin Health Solutions as applicable (referred to individually as “Company” and collectively as “Companies”).

APPLIES TO:

All lines of business except Medicare (*unless otherwise directed by a Medicare medical policy. Note that investigational services are considered “not medically necessary” for Medicare members.*)

BENEFIT APPLICATION

Medicaid Members

Oregon: Services requested for Oregon Health Plan (OHP) members follow the OHP Prioritized List and Oregon Administrative Rules (OARs) as the primary resource for coverage determinations. Medical policy criteria below may be applied when there are no criteria available in the OARs and the OHP Prioritized List.

DOCUMENTATION REQUIREMENTS

In order to determine the medical necessity of the request, the following documentation must be provided at the time of the request. All medical records and chart notes to include clinical documentation to support review including but not limited to the following:

- History
- Physical examination
- Treatment plan including whether surgery will be performed as multiple stages.
- If two (or more) surgeries are planned, each must be individually documented. This may (sometimes, but not necessarily) require multiple sets of photographs.

MEDICAL POLICY

Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)

- Color photographs with detail as described in [Policy Guidelines](#).
- If both a blepharoplasty and a ptosis repair are planned, both must be individually documented
- Detailed description of all “physical signs” when relevant to criteria. See Patient Complaints and Physical Signs in [Policy Guidelines](#).
- If planned procedure is reconstructive in circumstances of congenital defects, developmental abnormalities, trauma, infection, tumors, documentation of patient complaint regarding impacts on their ability to perform tasks of daily living (or, in the absence of a specific complaint, a statement that the repair is needed to prevent anticipated future damage to ocular structures).

POLICY CRITERIA

Note: This policy does not address blepharoplasty, blepharoptosis, or brow lifts requested as gender affirming interventions. Please see the [Policy Cross References](#) Section below for the appropriate policy.

- I. Blepharoplasty, blepharoptosis repair, and brow ptosis repair (brow lift) may be considered **medically necessary** to improve abnormal function when either of the following criteria are met (A. or B.):
 - A. Overhanging skin or upper lid position secondary to dermatochalasis, blepharochalasis, blepharoptosis, or pseudoptosis is sufficiently low to produce a visually-significant field restriction (approximately 30 degrees or less from fixation or Margin Reflex Distance of 2.0 mm or less) (see [Policy Guidelines](#)); **or**
 - B. Planned procedure is reconstructive (aims to improve abnormal function or approximate a normal appearance) in circumstances of congenital defects, developmental abnormalities, trauma, infection, tumors related to **any** of the following (note that documentation of visual field loss is not required – see [Policy Guidelines](#)):
 1. Ectropion, entropion or trichiasis,
 2. Repairs to address ocular exposure,
 3. Repairs to address difficulty fitting an ocular prosthesis,
 4. Primary essential idiopathic blepharospasm (uncontrollable spasms of the periorbital muscles) that is debilitating for which all other treatments have failed or are contraindicated,
 5. Prompt repair of an accidental injury or trauma.

Note: For patients with equal innervation to both eyelids, if one eyelid meets criterion I. above, the less ptotic eyelid may also be repaired.

- II. Blepharoplasty, blepharoptosis repair, and brow ptosis repair is considered **cosmetic and is not covered** when the above criterion I. is not met, including but not limited to when

MEDICAL POLICY

Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)

performed to enhance appearance in the absence of any signs and/or symptoms of functional abnormalities.

Link to [Policy Summary](#)

POLICY GUIDELINES

This policy may be primarily based on the following Center for Medicare and Medicaid Services (CMS) guidances:

- Local Coverage Determination (LCD): Blepharoplasty, Eyelid Surgery, and Brow Lift (L36286).¹

Lower Eyelid Blepharoplasty

Lower eyelid blepharoplasty is almost never functional in nature and is reviewed per policy criteria as written, though usually does not meet criteria.

Patient Complaints and Physical Signs

- A functional deficit or disturbance secondary to eyelid and/or brow abnormalities must be documented, such as interference with vision or visual field that impacts an activity of daily living.
- In addition, the documentation should show that the eye being considered for surgery has physical signs consistent with the functional deficit or abnormality.
- For Blepharoptosis Repair:
 - A margin reflex distance (MRD sometimes referred to as MRD1) of 2.0 mm or less. The MRD is a measurement from the corneal light reflex to the upper eyelid margin (NOT any overhanging skin that may be present causing pseudoptosis) with the brows relaxed, and
 - If applicable, the presence of Herring's effect (related to equal innervation to both upper eyelids) defying bilateral surgery when only the more ptotic eye clearly meets the MRD criteria (i.e., if lifting the more ptotic lid with tape or by instillation of phenylephrine drops into the superior fornix causes the less ptotic lid to drop downward and meet the strict criteria, the less ptotic lid is also a candidate for surgical correction.
- For Upper Blepharoplasty and/or Brow Ptosis Repair:
 - Redundant eyelid tissue hanging over the eyelid margin resulting in pseudoptosis where the "pseudo" margin produces a central "pseudo-MRD" of 2.0 mm or less, or
 - Redundant eyelid tissue predominantly medially or laterally that clearly obscures the line of sight in corresponding gaze.
- If an anatomic abnormality of the eye (such as an eccentric or elongated pupil) makes the MRD either difficult to establish or meaningless for this purpose, it is expected the surgeon will include a statement outlining their rationale that an equivalent standard has been met.

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

- Visual field loss must correlate with photographic appearance. Peripheral visual field testing performed with the eyelid and brow in the resting position (un-taped) and with eyelid and brow elevation (taped). Visual fields are not required for children (17 years or younger) or other patients physically or mentally unable to perform visual field testing (e.g. intellectual disability, severe neurologic disease).

Photographs

- Photographs are required to support upper eyelid surgery as medically necessary and covered.
- The “physical signs” in the Documentation Requirements and Policy Guidelines must be clearly represented in photographs of the structures of interest and the photographs must be of good quality and of sufficient size and detail as to make those structures easily recognizable.
- The patient’s head and the camera must be in parallel planes, not tilted so as not to distort the appearance of any relevant finding (e.g., a downward head tilt might artificially reduce the apparent measurement of a MRD).
- Unless medial/lateral gaze is required to demonstrate a specific deficit, photos should be with gaze in the primary position, looking straight ahead.
- Oblique photos are only necessary if needed to better demonstrate a finding not clearly shown by other requested photos.
- Digital or film photographs are acceptable, and may be submitted electronically where possible. Photographs must be identified with the member’s name and the date.
- For Blepharoptosis Repair:
 - Photographs of both eyelids in the frontal (straight-ahead) position should demonstrate the MRD outlined in Policy Guidelines. If the eyelid obstructs the pupil, there is a clear-cut indication for surgery. (For reference, the colored part of the eye is about 11 mm in diameter, so the distance between the light reflex and the lid would need to be about one fifth that distance or less for the MRD to be 2.0 mm or less.)
 - In the special case of documenting the need for bilateral surgery because of Herring’s law, two photos are needed:
 - One showing both eyes of the patient at rest demonstrating the above MRD criterion in the more ptotic eye, and
 - Another showing both eyes of the patient with the more ptotic eyelid raised to a height restoring a normal visual field, resulting in increased ptosis (meeting the above MRD standard) in the less ptotic eye.
- Reviewers will assume the accepted average of 11 mm of corneal diameter to assess measurements in photographs. If a patient’s corneal diameter deviates from this by more than 0.5 mm, this should be clearly documented in the record so appropriate adjustments can be made. Alternatively, an accurate millimeter rule can be taped along the brow, on the cheek, or elsewhere in the photo (approximately in the corneal plane) to facilitate such measurements.
- For Upper Lid Blepharoplasty:
 - Photographs of the affected eyelid(s) in both frontal (straight ahead) and lateral (from the side) positions demonstrate the physical signs in Policy Guidelines. Oblique photos

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

are only necessary if needed to better demonstrate a finding not clearly shown by frontal and lateral photos.

- For Brow Ptosis Repair:
 - One frontal (straight ahead) photograph should document drooping of a brow or brows and the appropriate other criteria in Policy Guidelines. If the goal of the procedure is improvement of dermatochalasis, a second photograph should document such improvement by manual elevation of brow(s). If a single frontal photograph that includes the brow(s) would render other structures too small to evaluate, additional (overlapping to the degree possible) photos should be taken of needed structures to ensure all required criteria can be reasonably demonstrated and evaluated.
- If both a blepharoplasty and a ptosis repair are planned, both must be individually documented. This may (sometimes, but not necessarily) require two sets of photographs: showing a pseudo-MRD of 2.0 mm or less secondary to the redundant skin (and its correction by taping), AND an MRD of 2.0 mm or less secondary to the blepharoptosis.

Definitions relevant to the policy criteria:

Activities of daily living: The activities of daily living (ADLs) is a term used to describe essential skills that are required to independently care for oneself. Examples may include, but are not limited to, the following:

- Ambulating
- Feeding
- Dressing
- Personal hygiene
- Transportation and shopping
- Meal preparation
- Housecleaning and home maintenance

Blepharoptosis: drooping of the upper eyelid related to the position of the eyelid margin with respect to the visual axis

Brow ptosis: drooping of the eyebrows to such an extent that excess tissue is pushed into the upper eyelid that may cause mechanical blepharoptosis and/or dermatochalasis

Blepharoplasty: removal of eyelid skin, fat, and or muscle

Blepharoptosis repair (also known as levator resection): restoring the eyelid margin to its normal anatomic position.

Brow ptosis repair: restoring the eyebrow tissues to their normal anatomic position.

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

Dermatochalasis: excess skin with loss of elasticity that is usually the result of the aging process.

Ectropion: an outward turning of the eyelid margin.

Entropion: an inward turning of the eyelid margin and appendages such that the pilosebaceous unit and mucocutaneous junction are directed posterior towards the globe.

Trichiasis: a condition, resulting from the eye infection called trachoma, in which the eyelid turns inward and eyelashes rub against the eye, resulting in corneal scarring and loss of vision.

BILLING GUIDELINES

When a Medically Necessary Procedure is Performed with a Cosmetic Procedure

When a cosmetic and noncovered surgical procedure is performed in the same operative session as a medically necessary and covered surgical procedure, medical necessity review will be conducted for the allowable procedure only. For example, if dermatochalasis would be resolved sufficiently by brow ptosis repair alone, an upper lid blepharoplasty in addition would be considered cosmetic. Similarly, if a visual field deficit would be resolved sufficiently by upper lid blepharoplasty alone (for tissue hanging over the lid margin), a blepharoptosis repair in addition would be considered cosmetic.

CPT CODES

All Lines of Business Except Medicare	
Prior Authorization Required	
15820	Blepharoplasty, lower eyelid
15821	Blepharoplasty, lower eyelid; with extensive herniated fat pad
15822	Blepharoplasty, upper eyelid
15823	Blepharoplasty, upper eyelid; with excessive skin weighting down lid
67900	Repair of brow ptosis (supraciliary, mid-forehead or coronal approach)
67901	Repair of blepharoptosis; frontalis muscle technique with suture or other material (eg, banked fascia)
67902	Repair of blepharoptosis; frontalis muscle technique with autologous fascial sling (includes obtaining fascia)
67903	Repair of blepharoptosis; (tarso) levator resection or advancement, internal approach
67904	Repair of blepharoptosis; (tarso) levator resection or advancement, external approach
67906	Repair of blepharoptosis; superior rectus technique with fascial sling (includes obtaining fascia)
67908	Repair of blepharoptosis; conjunctivo-tarso-Muller's muscle-levator resection (eg, Fasanella-Servat type)

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

No Prior Authorization Required	
67909	Reduction of overcorrection of ptosis
67911	Correction of lid retraction
67914	Repair of ectropion; suture
67915	Repair of ectropion; thermocauterization
67916	Repair of ectropion; excision tarsal wedge
67917	Repair of ectropion; extensive (eg, tarsal strip operations)
67921	Repair of entropion; suture
67922	Repair of entropion; thermocauterization
67923	Repair of entropion; excision tarsal wedge
67924	Repair of entropion; extensive (eg, tarsal strip or capsulopalpebral fascia repairs operation)
67930	Suture of recent wound, eyelid, involving lid margin, tarsus, and/or palpebral conjunctiva direct closure; partial thickness
67935	Suture of recent wound, eyelid, involving lid margin, tarsus, and/or palpebral conjunctiva direct closure; full thickness

DESCRIPTION

Blepharoplasty and blepharoptosis repair may be defined as any eyelid surgery that improves abnormal function, reconstructs deformities, or enhances appearance. They may be either functional/reconstructive or cosmetic. Upper blepharoplasty (removal of upper eyelid skin) and/or repair of blepharoptosis should be considered functional/reconstructive in nature when the upper lid position or overhanging skin is sufficiently low to produce functional complaints, usually related to visual field impairment whether in primary gaze or down-gaze reading position. Upper blepharoplasty may also be indicated for chronic dermatitis due to redundant skin. Another indication for blepharoptosis surgery is patients with an anophthalmic socket experiencing ptosis or prosthesis difficulties. A brow lift is a surgery that may be performed to correct brow ptosis (droop of the eyebrows), which can also produce or contribute to functional impairment.

Primary essential idiopathic blepharospasm is characterized by severe squinting, secondary to uncontrollable spasms of the periorbital muscles. Occasionally, it can be debilitating if other treatments have failed or are contraindicated. In these rare cases, an extended blepharoplasty with wide resection of the orbicularis oculi muscle complex may be necessary to properly treat symptoms.

CLINICAL PRACTICE GUIDELINES

American Society of Ophthalmic Plastic and Reconstructive Surgery (ASOPRS)

The 2015 ASOPRS guidelines on “Functional Blepharoplasty, Blepharoptosis, and Brow Ptosis Repair”² recommend the following:

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

“Recommended coverage indications

- I. Upper eyelid blepharoplasty (CPT 15823) should be considered medically necessary when documentation demonstrates:
 - A. The patient’s complaint of interference with daily visual tasks or visual field-related activities, and
 - B. Visual obstruction due to excessive overhanging skin resting on or depressing the lashes or eyelid margin. Visual obstruction is defined by peripheral visual field testing consistent with the recommended documentation requirement (see below).

- II. Upper eyelid ptosis repair (CPT 67901, 67902, 67903, 67904, 67906, 67908) should be considered medically necessary when documentation demonstrates:
 - A. The patient’s complaint of interference with vision or visual field-related activities, and
 - B. A margin to reflex distance (MRD) less than or equal to 2 mm in primary or downgaze.
 - C. Visual obstruction due to ptotic upper eyelid. Visual obstruction is defined by peripheral visual field testing consistent with the recommended documentation requirement (see below).
 - D. The position of one upper eyelid, which initially appears not to meet criteria but becomes more ptotic with an MRD of 2 mm or less when the other, more ptotic eyelid is elevated (i.e. Hering’s Law).

- III. Repair of brow ptosis (CPT 67900) should be considered medically necessary when documentation demonstrates:
 - E. Brow ptosis to the extent it contributes to skin fold overlap and/or blepharoptosis meeting the criteria outlined above for upper eyelid blepharoplasty and/or ptosis surgery.

Recommended documentation requirements

1. Clinical notes documenting patient complaints of visual impairment secondary to abnormal eyelid or brow position resulting in limitation of daily activities such as reading, driving, and difficulty seeing objects approaching from the periphery, or redundant upper eyelid skin resulting in looking through the eyelashes or seeing the upper eyelid skin.
2. Clinical notes documenting an MRD of 2 mm or less for blepharoptosis repair.
3. Peripheral visual field testing performed with the eyelid and brow in the resting position that documents a baseline superior visual field of 30 degrees or less from fixation and improvement of at least 12 degrees over baseline with eyelid and brow elevation. Both manual, including Goldmann and tangent screen, and automated fields are acceptable. The measurement of a patient’s superior visual field is defined as the lowest point seen at the vertical meridian. Visual fields are not required for children or other patients physically unable to perform visual field testing.
4. Photographs documenting the above including at least a good quality frontal upper face photograph, with the gaze in primary position clearly showing the corneal light reflex unless

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

- the ptotic tissue is so severe as to obscure the light reflex thus demonstrating severe ptosis and/or dermatochalasis.
5. Oblique or lateral photographs if helpful in further demonstrating overhanging excess skin as well as lash ptosis due to mechanical displacement by the overhanging skin fold.
 6. Frontal photograph with the patient looking in down-gaze documenting those cases in which the ptosis is worse in the down-gaze position.
 7. Photograph with the brows elevated or taped up to a normal position to document the effect of brow ptosis when both eyelid ptosis repair and brow ptosis repair are planned.”

American Academy of Ophthalmology (AAO)

The 2011 AAO position statement on “Functional Indications for Upper Eyelid Ptosis and Blepharoplasty Surgery”³ stated the following:

“Ptosis and upper eyelid blepharoplasty surgery were found to be functionally beneficial for each of these quantitative findings:

- MRD₁ of ≤ 2 mm measured in primary gaze
- Superior visual field loss of 12 degrees or 24%
- Down-gaze ptosis impairing reading documented by MRD₁ of ≤ 2 mm⁷ measured in down gaze

Ptosis and upper eyelid blepharoplasty were also found to be functionally beneficial for the following qualitative findings:

- Self-reported functional impairment from upper eyelid droop
- Chin-up backward head tilt induced by visual field impairment caused by lids
- Interference with occupational duties and safety resulting from visual impairment caused by the upper lids
- Symptoms of discomfort, eye strain, or visual interference due to the upper eyelid position

The reviewed literature did not provide strong data on the following functional indications for ptosis and blepharoplasty surgery:

- Dermatitis
- Difficulty wearing a prosthesis in an anophthalmic socket
- Temporal visual field impairment preventing a driver from meeting licensing standards

Ptosis and dermatochalasis can occur concomitantly. Each has its own functional indications for repair, and different surgical procedures are required to correct them.”

The 2017 AAO Amblyopia Preferred Practice Pattern clinical practice guidelines on amblyopia, stated the following:⁴

- “All children with amblyopia should be offered treatment regardless of age.

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	--

- Surgery to treat the cause of amblyopia may be indicated when the cause of the amblyopia can be attributed to blepharoptosis that is severe enough to prevent successful amblyopia therapy without surgical correction.”

POLICY SUMMARY

The medical necessity of blepharoplasty, blepharoptosis and brow lift procedures for the various indications addressed in this policy are primarily based on clinical rationale and current evidence-based clinical practice guidelines. When medical necessity criteria for these procedures are not met, these procedures are considered to be cosmetic in nature.

INSTRUCTIONS FOR USE

Company Medical Policies serve as guidance for the administration of plan benefits. Medical policies do not constitute medical advice nor a guarantee of coverage. Company Medical Policies are reviewed annually and are based upon published, peer-reviewed scientific evidence and evidence-based clinical practice guidelines that are available as of the last policy update. The Companies reserve the right to determine the application of Medical Policies and make revisions to Medical Policies at any time. Providers will be given at least 60-days’ notice of policy changes that are restrictive in nature.

The scope and availability of all plan benefits are determined in accordance with the applicable coverage agreement. Any conflict or variance between the terms of the coverage agreement and Company Medical Policy will be resolved in favor of the coverage agreement.

REGULATORY STATUS

Mental Health Parity Statement

Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case. In cases where medical necessity is not established by policy for specific treatment modalities, evidence not previously considered regarding the efficacy of the modality that is presented shall be given consideration to determine if the policy represents current standards of care.

MEDICAL POLICY CROSS REFERENCES

- Cosmetic and Reconstructive Surgery
- Gender Affirming Interventions

MEDICAL POLICY	Eye: Blepharoplasty, Blepharoptosis Repair, and Brow Lift (All Lines of Business Except Medicare)
-----------------------	---

REFERENCES

1. Centers for Medicare & Medicaid Services. Local Coverage Determination (LCD): Blepharoplasty, Eyelid Surgery, and Brow Lift (L36286). <https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=36286>. Accessed 6/9/2022.
2. American Society of Ophthalmic Plastic and Reconstructive Surgery (ASOPRS). White Paper on Functional Blepharoplasty, Blepharoptosis, and Brow Ptosis Repair. Approved: 01/15/2015. <https://www.asoprs.org/assets/docs/1%20-%20FINAL%20ASOPRS%20White%20Paper%20January%202015.pdf>. Accessed 6/9/2022.
3. Cahill KV, Bradley EA, Meyer DR, et al. Functional indications for upper eyelid ptosis and blepharoplasty surgery: a report by the American Academy of Ophthalmology. *Ophthalmology*. 2011;118(12):2510-2517. <https://www.ncbi.nlm.nih.gov/pubmed/22019388>
4. American Academy of Ophthalmology. Amblyopia Preferred Practice Pattern. Pediatric Ophthalmology/Strabismus Panel. 2017. <https://www.aao.org/preferred-practice-pattern/amblyopia-ppp-2017>. Accessed 6/9/2022.