

Spinal Stabilization Devices and Interspinous Spacers

MEDICAL POLICY NUMBER: 19

Effective Date: 6/1/2025	COVERAGE CRITERIA	2
Last Review Date: 5/2025	POLICY CROSS REFERENCES	3
Next Annual Review: 5/2026	POLICY GUIDELINES	3
	REGULATORY STATUS.....	5
	CLINICAL EVIDENCE AND LITERATURE REVIEW.....	6
	HEALTH EQUITY CONSIDERATIONS	13
	BILLING GUIDELINES AND CODING	13
	REFERENCES	14
	POLICY REVISION HISTORY	16

INSTRUCTIONS FOR USE: Company Medical Policies serve as guidance for the administration of plan benefits. Medical policies do not constitute medical advice nor a guarantee of coverage. Company Medical Policies are reviewed annually and are based upon published, peer-reviewed scientific evidence and evidence-based clinical practice guidelines that are available as of the last policy update. The Company reserves the right to determine the application of medical policies and make revisions to medical policies at any time. The scope and availability of all plan benefits are determined in accordance with the applicable coverage agreement. Any conflict or variance between the terms of the coverage agreement and Company Medical Policy will be resolved in favor of the coverage agreement. Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case. In cases where medical necessity is not established by policy for specific treatment modalities, evidence not previously considered regarding the efficacy of the modality that is presented shall be given consideration to determine if the policy represents current standards of care.

SCOPE: Providence Health Plan, Providence Health Assurance and Providence Plan Partners as applicable (referred to individually as “Company” and collectively as “Companies”).

PLAN PRODUCT AND BENEFIT APPLICATION

☒ Commercial

☐ Medicaid/OHP*

☐ Medicare**

*Medicaid/OHP Members

Oregon: Services requested for Oregon Health Plan (OHP) members follow the OHP Prioritized List and Oregon Administrative Rules (OARs) as the primary resource for coverage determinations. Medical policy criteria below may be applied when there are no criteria available in the OARs and the OHP Prioritized List.

Spinal Stabilization Devices and Interspinous Spacers – Guideline Note 173

**Medicare Members

This Company policy may be applied to Medicare Plan members only when directed by a separate Medicare policy. Note that investigational services are considered “**not medically necessary**” for Medicare members.

COVERAGE CRITERIA

Note: The iFuse Implant System™ for sacroiliac joint fusion is addressed in the Medical Policy, [Sacroiliac Joint Fusion or Stabilization](#).

Dynamic Stabilization Devices

- I. The dynamic stabilization devices listed below (228999, 64999, L8699) are considered **not medically necessary**, including, but not limited to, the following:
 - A. Aspen Spinous Fixation System
 - B. AccuFlex™ System
 - C. BioFlex System
 - D. Bronsard’s Ligament
 - E. CD Horizon Agile™ Dynamic Stabilization Devices
 - F. CD Horizon Spire Fixation System
 - G. Cosmic™ Posterior Dynamic System
 - H. DSS Stabilization
 - I. DSS (Dynamic Soft Stabilization) System
 - J. DTO (Dynesys-to-Optima)
 - K. Dynabolt™ Dynamic Stabilization System
 - L. Dynesys®
 - M. Expedium™
 - N. FASS (Fulcrum Assisted Soft Stabilization)
 - O. Graf Ligament

- P. IsoBar® Spinal System
- Q. Leeds-Keio Ligamentoplasty
- R. LemiFlex Spinal Stabilization
- S. NFix™ II Dynamic Stabilization
- T. NFlex™ Controlled Motion System
- U. REVERE Stabilization System
- V. Satellite™ Spinal System
- W. Stabilimax NZ Dynamic Spine Stabilization System
- X. TRANSITION® Stabilization System
- Y. Viper™
- Z. Zodiac DynaMo System

Interspinous Spacers

- II. The interspinous spacers listed below (22867, 22868, 22869, 22870, C1821) are considered **not medically necessary**, including, but not limited to, the following:
 - A. Coflex® Interlaminar Stabilization Device
 - B. Superior® Interspinous Spacer System
 - C. Aperius™ – PercLID™ System
 - D. DIAM™ Spinal Stabilization System
 - E. Falena® Interspinous Decompression Device
 - F. FLEXUS™
 - G. Helifix® Interspinous Spacer System
 - H. In-Space
 - I. NL-Prow™ Interspinous Spacer
 - J. Stenofix
 - K. Wallis System®

Link to [Evidence Summary](#)

POLICY CROSS REFERENCES

- [Sacroiliac Joint Fusion or Stabilization \(Company\)](#), MP24

The full Company portfolio of current Medical Policies is available online and can be [accessed here](#).

POLICY GUIDELINES

BACKGROUND

Lumbar Spinal Stenosis (LSS)

LSS is predominantly caused by degeneration in the intervertebral discs, ligaments and bone structures of the spine, and is characterized by a narrowing of the spinal canal, lateral spinal recesses and compressed neural elements in the lower back, resulting in pain and disability. Symptoms are typically provoked by upright exercise, including walking, and relieved with forward flexion at the waist, sitting or reclining. While conservative treatments improve symptoms in one-third of patients (e.g. physical therapy, nonsteroidal anti-inflammatory drugs), surgical options for those refractory to these therapies range from minimally invasive decompression techniques to traditional surgical laminectomy with or without spinal fusion, laminotomy, or hemilaminotomy.

Dynamic Stabilization Devices

Dynamic stabilization devices provide an adjunct or alternative to spinal fusion for the treatment of chronic instabilities or deformities of the thoracic, lumbar, and sacral spine, including, but not limited to, degenerative spondylolisthesis or previous failed spinal fusion. In contrast to rigid devices that fully stabilize affected spinal segments, dynamic stabilization devices use flexible materials – anchored to the vertebrae by either synthetic cords or pedicle screws – which purport to preserve some measure of mobility of the spinal segment while also stabilizing the joint.

Aspen Spinous Fixation System

The Aspen Spinous Fixation System is designed for posterior fixation to promote fusion from T1 to S1 vertebrae.¹ The System consists of a “spiked plate” implant that fixates the spine in single- or multi-level constructs, supporting the formation of fusion and decompression by fixation, load sharing and interspinous process spacing. The device is used as an adjunct to interbody fusion and/or posterior fusion with decompression.

DSS Stabilization System

The DSS Stabilization System is a single-level, pedicle-based fixation system from the T4 to S1 vertebrae, consisting of polyaxial, cannulated pedicle screws, slotted couplers, and rigid couplers. The System purports to immobilize and stabilize spinal segments as an adjunct to fusion in the treatment of acute and chronic instabilities or deformities of the thoracic, lumbar, and sacral spine.²

Dynesys®

Dynesys is a semi-rigid pedicle-based dynamic stabilization system consisting of three components: titanium screws anchor the system to the spine; polycarbonate urethane spacers limit spinal extension; and polymer cords that act as a tension band to limit spinal flexion.³ The device theoretically preserves motion of the treated segment while also alleviating pain by restricting loading on adjacent discs and facet joints.

Isobar Spinal System

Isobar is a semi-rigid, pedicle-based system intended to immobilize and stabilize spinal segments as an adjunct to fusion, for the treatment of spondylolisthesis.

Interspinous Spacers

Interspinous spacers are small devices, implanted between vertebral spinous processes at one or two vertebral levels, that stabilize or distract adjacent lamina and/or spinous processes. The spacers are thought to alleviate pain in patients with spinal stenosis and neurogenic claudication by expanding the neural foramen, decompressing the nerves and limiting painful lumbar extension, while maintaining the flexion of the spinal interspace.

Coflex® Interlaminar Stabilization Device

The Coflex Interlaminar Stabilization Device is a U-shaped titanium alloy implant that is inserted after decompression of stenosis.⁴ The device is positioned horizontally and is appropriate for 1 level or 2 contiguous levels from lumbar (L) vertebrae L1 to L5. The device purports to alleviate pain by limiting lumbar spinal extension, unloading facet joints and stabilizing the motion segment at the treated vertebral level(s).

Superion Interspinous Spacer System (ISS)

The Vertiflex™ Superion Interspinous Spacer (ISS) (Boston Scientific Corporation) first attained FDA premarket approval (PMA) in 2015 (P140004), and was the predicate device to what is now marketed as the Superion™ Indirect Decompression System (IDS).⁵ The Vertiflex™ Superion™ IDS is a minimally invasive treatment option for lumbar spinal stenosis (LSS).⁶ No surgical decompression is used in this surgery. The system is indicated for skeletally mature patients experiencing pain, numbness, and/or cramping in the legs (neurogenic intermittent claudication) secondary to a diagnosis of moderate degenerative lumbar spinal stenosis, with or without grade 1 spondylolisthesis, confirmed by X-ray, MRI, and/or CT evidence of thickened ligamentum flavum, narrowed lateral recess, and/or central canal or foraminal narrowing. It is intended for patients with impaired physical function who experience relief in flexion from symptoms of leg/buttock/groin pain, numbness, and/or cramping, with or without back pain, and who have undergone at least six months of nonoperative treatment. FDA PMA Product Code: NQO.

The Vertiflex™ Superion is one of only two interspinous spacer or spinous process plates with full FDA-approval. Following worldwide recalls, the formerly approved X-STOP devices had FDA-approval formally withdrawn in 2015.⁷

REGULATORY STATUS

U.S. FOOD AND DRUG ADMINISTRATION (FDA)

Approval or clearance by the Food and Drug Administration (FDA) does not in itself establish medical necessity or serve as a basis for coverage. Therefore, this section is provided for informational purposes only.

The following dynamic stabilization devices and interspinous spacers have received FDA clearance:

- Aspen Spinous Fixation System⁸

- DSS Stabilization System⁹
- Dynesys^{®10}
- Isobar Spinal System¹¹
- Coflex[®] Interlaminar Stabilization Device¹²
- Superion Interspinous Spacer System⁵

CLINICAL EVIDENCE AND LITERATURE REVIEW

EVIDENCE REVIEW

Dynamic Stabilization Devices

Aspen Spinous Fixation System

Evidence for the Aspen Spinous Fixation System is limited to two cadaveric studies describing the device's biomechanical effects.^{13,14} No studies published within the past five years were identified.

DSS Stabilization System

In 2018, Bieri and colleagues conducted a retrospective cohort study of clinical outcomes for dynamic posterior stabilization.¹⁵ Patients (mean age: 67 years) were treated with either the DSS Stabilization System (n=202) or underwent posterior lumbar intervertebral fusion (PLIF) (n=269). Median follow-up was 3.3 years. The primary outcome of interest was the change in the patient-reported Core Outcome Measures Index score (COMI; a 0-10 scale). Investigators reported no difference between groups in improved COMI score, back pain relief, leg pain relief, blood loss and complications. Compared to the PLIF group, patients treated with DSS experienced longer hospital stays ($p = 0.03$) but fewer repeat surgeries ($p = 0.01$), and shorter surgery times ($p < 0.001$). Investigators concluded that while DSS may be a viable alternative to PLIF at mid-term follow-up, multicenter studies were needed to confirm reported findings. Limitations include the study's retrospective design, heterogeneity of patient characteristics between groups at baseline, potential underreporting of complication rates, and the lead author's financial conflict of interest with the manufacturer of the DSS Stabilization System.

Dynesys[®]

In 2016, Lee and colleagues conducted a meta-analysis evaluated the efficacy of the Dynesys system versus posterior lumbar interbody fusion (PLIF) for the treatment of degenerative lumbar spinal disease.¹⁶ In total, 7 studies (n=506) (1 RCT, 2 prospective cohort studies and 4 retrospective cohort studies), all of which compared clinical and radiological outcomes, were included for review. Average follow-up was 2 years. Pooled analysis found no significant difference in disability, back pain, leg pain, complication rates and length of hospital stay between the two groups. Limitations include differences in treatment parameters across studies, and the low-middle income treatment context (i.e. China) for 4 out of 7 studies. No eligible studies were conducted in the United States.

Isobar Spinal System

- In 2016, two retrospective studies in China (n=37) evaluated the efficacy of the Isobar Spinal System.^{17,18} Follow-up ranged from 12 to 53 months. Both studies reported significant pain relief

compared to baseline. Investigators of both studies called for larger, prospective studies with longer follow-up to better establish Isobar's safety and efficacy.

- In 2014, Fu and colleagues prospectively evaluated the functional and radiological outcomes of Isobar in conjunction with spinal fusion.¹⁹ In total, 36 patients underwent posterior Isobar dynamic stabilization for single-level degenerative lumbar disc disease with instability and mild adjacent level degeneration. At 24-months follow-up, patients experienced significant improvement compared to baseline in both mean visual analog scale scores, and Oswestry Disability Index scores. While intervertebral angle (IVA) at the adjacent level increased, disc height at the index and adjacent levels and intervertebral angle (IVA) at the index level decreased (i.e. degenerated) at each follow-up, No adverse events or reoperations were reported. Limitations include the study's small sample size, inadequate follow-up and lack of comparator groups. Investigators concluded that additional studies with long-term follow-up were necessary to establish Isobar's safety and efficacy.

Interspinous Spacers

Coflex® Interlaminar Stabilization Device

- In 2022, Hayes updated a health technology assessment evaluating the safety and efficacy of Coflex Interlaminar Stabilization Device for the treatment of lumbar spinal stenosis (LSS).²⁰ Hayes systematically searched the literature through August 2018, identified eligible studies, assessed quality and extracted data. In total 12 publications were included for review, each of which compared treatments with more than 50 patients for at least 12-months. Sample sizes ranged from 62 to 254 patients. Outcomes of interest included disability, pain, function, quality of life, radiographic outcomes of range of motion and stability; secondary surgical interventions; procedural outcomes and safety.

Across 9 studies, Coflex was generally associated with clinically important improvements in patient disability, with few significant between-group differences in 5 studies comparing Coflex with fusion at ≤60 month follow-up. Among 2 studies comparing Coflex with decompression alone at 2-year follow-up, no significant differences were reported. Coflex was associated with significant improvements in pain (assessed by visual analog score) from baseline at 60-months. Studies reported no significant differences between Coflex and fusion or decompression alone. Studies reported improvements in functional outcomes, quality of life, narcotic use, secondary surgical interventions and safety among both Coflex and comparator groups. Compared to fusion and decompression, Coflex was favored for both radiographic outcomes of range of motion and stability, and procedural outcomes.

Hayes judged the overall evidence base evaluating Coflex to be of low-quality and assigned a "C" rating (potential but unproven benefit). Results indicated that Coflex was associated with improvements in pain, function and disability comparable to fusion or decompression alone, yet results' validity was limited by heterogeneous patient populations, small sample sizes, a lack of randomization, inadequate follow-up, inadequate blinding, and the lack of patient selection criteria. Hayes concluded that there was insufficient evidence to draw conclusions regarding the use of the Coflex device for the treatment of LSS.

- In 2018, Mo and colleagues conducted a network meta-analysis and systematic review evaluating the efficacy and safety of posterior lumbar interbody fusion compared to interspinous process devices (IPD), including Coflex.²¹ Investigators systematically searched appropriate databases, identified relevant studies, assessed quality, extracted data and pooled results. In total, 27 studies (n=2,241) were included for review. Investigators found no significant differences between groups in Oswestry Disability Index, visual analogue scale, Japanese Orthopedic Association Scores, and posterior disc height. Patients receiving IPDs experienced significantly more range of motion and less adjacent segment degeneration, but no improvements in pain relief, quality of life, disc space height and lumbar function. Limitations include the low-middle income treatment context of 24 of the included 27 studies (i.e. China), which may limit results' generalizability.
- In 2017, Zhao and colleagues conducted a systematic review and meta-analysis evaluating the efficacy of Coflex and other interspinous process devices (IPD) alone versus bony decompression surgery for the treatment of lumbar spinal stenosis (LSS).²² Investigators systematically searched appropriate databases, identified relevant studies, assessed quality, extracted data and pooled results. In total, 7 publications deriving from 4 high-quality RCTs were included for review (n= 400). Follow-up was ≤24 months. Investigators reported no significant difference between the two groups in length of hospital stay, visual analog score (VAS) leg pain scores, and complication rates, but higher VAS low back pain scores and reoperation rates for IPD patients. Two studies demonstrated inferior quality of life improvements relative to bony decompression. Study limitations included the variety of IPD devices used across trials, inadequate follow-up among individual RCTs and the inability of investigators to conduct sub-group analyses given the lack of relevant data. Investigators concluded that while both IPD and bony decompression were "acceptable treatment strategies" for LSS, results did not indicate IPD's superiority to bony decompression – current gold standard treatment. Investigators called for larger studies with longer follow-up to better establish the safety and efficacy of IPDs.
- In 2017, Machado and colleagues conducted a Cochrane review to evaluate the efficacy of surgery in the management of patients with lumbar spinal stenosis (LSS) and the comparative efficacy between commonly performed surgical techniques.²³ Investigators searched the literature through June 2016 for RCTs that compared efficacy and safety of surgery compared with no treatment, placebo or sham surgery, or with another surgical technique. Follow-up ranged from 1 to 2 years. Outcomes of interest were pain intensity, physical function or disability status, quality of life and recovery. Secondary outcomes included measurements related to surgery (e.g. perioperative blood loss, length of hospital stay, reoperation rates).

In total, 39 RCTs were included for review (n=2352), all of which compared two or more surgical techniques. All trials were assessed to be at high risk of bias due to inadequate blinding, incomplete randomization and a lack of intention-to-treat analyses. Overall, investigators reported no differences for primary and secondary outcomes. Three trials compared interspinous process spacer devices, including Coflex, to conventional bony decompression. Spacers achieved similar reductions in pain, disability, and perioperative blood loss, but longer operation times and incurred higher risks of reoperation. Two trials compared interspinous spacer devices, including Coflex, with decompression plus fusion. Among patients receiving spacer devices, investigators found no difference in pain relief or rate of reoperations, but significant improvements in disability reduction, operation time and perioperative blood loss. Investigators concluded that evidence is insufficient to recommend interspinous process spacers over decompression alone.

- In 2015, Moojen and colleagues published results from their double-blind RCT evaluating the efficacy of interspinous process devices (IPDs), including Coflex, versus conventional surgical decompression for lumbar spinal stenosis.²⁴ In total, 211 participants were randomized to one of two groups – IPD (n=80) and spinal bony compression (n=79) – and treated at one of five treatment centers across The Netherlands. At 2-year follow-up, investigators found no significant differences in success rates between groups, but significant increases in reoperations and visual analog scale back pain among the IPD group. Investigators concluded that these increases in the IPD group “suggest inferiority [of IPD] compared to classical decompression.”

Superion Interspinous Spacer System (ISS)

- In January 2022, ECRI published an updated clinical evidence assessment of the Superion IDS for treating lumbar spinal stenosis.²⁵ The ECRI authors evaluated full-text of 10 studies reporting on over 4,500 unique patients published between January 1, 2010 and July 6, 2022. Their findings included:
 - 1 RCT (Patel et al. 2015, n = 391) compared Superion (n = 190) with X-Stop (n = 201) implantation in patients with moderate LSS and reported on Zurich Claudication Questionnaire (ZCQ) domains, reoperation rates, and AEs through 36-month follow-up. This study was the Superion Investigational Device Exemption (IDE) trial.
 - 1 retrospective, multicenter chart review (Hagedorn et al. 2011, n = 199) compared Superion (n = 124) with the minimally invasive lumbar decompression (MILD) procedure (n = 57) in patients with LSS and reported on the rate of lumbar decompression revisions through 2-year follow-up.
 - 1 retrospective chart review (Welton et al. 2021, n = 567) compared Superion (n = 189) with laminectomy or laminotomy (n = 378) in patients with symptomatic LSS and reported on AEs through 30-day follow-up and device-specific AEs for Superion through 2-year follow-up.
 - 1 multicenter, historical control study (Nunley et al. 2017, n = 597) compared data from the Superion arm of the Superion IDE trial (n = 190) with decompressive laminectomy data from 2 published studies (n = 407) in patients with moderate degenerative LSS and reported on effect sizes of Oswestry Disability Index (ODI) scores at 2-, 3-, and 4-year follow-up and 2-year ZCQ symptom severity and physical function domains.
 - 1 multicenter, historical control study (Lauryssen et al. 2015, n = 1,235) compared data from the Superion arm of the Superion IDE trial (n = 190) with decompressive laminectomy (using data from 19 published studies [n = 1,045]) in patients with LSS and reported on ODI score improvement at 24-month follow-up, visual analog scale (VAS) leg and back pain, and ZCQ symptom severity and physical function domains.
 - 1 retrospective, multicenter before-and-after study (Nunley et al 2018, n = 189) compared Short Form (SF-12) physical and mental component scores before and 5 years after Superion implantation. Study used data from the Superion arm of the Superion IDE trial data.
 - 1 prospective, single-center before-and-after study (Bini et al. 2011, n = 121) compared ODI scores, extremity and axial pain scores (0 to 10 scale), and health-related quality of life (SF-36) physical and mental component scores before and 12 months after Superion implantation and reported clinical success rates and reported AEs through 12-month follow-up.

- 1 prospective multicenter registry (Tekmyster et al. 2019, n = 2,090) assessed Superion implantation in patients with symptomatic LSS and reported on improvements in mean leg pain severity, back pain severity, patient satisfaction, and revision rate through 12-month follow-up.
- 1 cost-effective study (Parker et al. 2015) compared Superion with decompressive surgery and conservative care in patients with LSS and reported on costs and quality-adjusted life-years.

ECRI reported that the evidence is “somewhat favorable” with the following limitations:

“The RCT was small and had high attrition at five-year follow-up. The nonrandomized comparative studies, historical control studies, and before-and-after studies are all at high risk of bias due to three or more of the following: single-center focus, small sample size, retrospective design, and lack of randomization and independent controls. Two studies (Laurysen et al. and Tekmyster et al.) did not report on statistical analyses for some patient outcomes. Only one study provided context for the reasons for reoperations; reoperation reasons (whether due to lack of effectiveness, device-related or procedure-related AEs) are unclear in the other studies, which limits interpretation of results and assessment of comparative AEs. Two historical control studies, two nonrandomized comparison studies, and two before-and-after studies assessed the same group of Superion-treated patients; thus, independent RCTs comparing Superion with other devices and laminectomy are needed to validate findings..”

- In 2023, Hayes published an evolving evidence review of full-text clinical studies evaluating the Superion Interspinous Spacer for the treatment of lumbar spinal stenosis with neurogenic claudication.²⁶ The authors identified “minimal support” for using the Superion ISS system. This level of support was reflected by their findings:
 - “Studies were of very poor or poor quality and no comparative studies were identified
 - Post-surgical improvements in pain and disability are reported at up to 3 years follow-up
 - Studies do not demonstrate equal or superior benefits or advantages over commercially available alternatives or fusion surgery”

The Hayes authors found conflicting information in professional society guidelines, noting that “several organizations recommend minimally invasive interspinous distraction procedures, in general, for carefully selected patients. This level of support reflects:

- Recommendations from guidelines are mixed, with some determining evidence to be sufficient to support use, while others determined evidence is insufficient to support use.
- One of the 2 guidelines finding sufficient evidence was published prior to the publication of any of the studies included in this report; the other is based on the Investigating Superion™ In Spinal Stenosis study

- No guidelines specifically naming the Superion Interspinous Spacer were identified, although the 2019 MIST guideline refers to a study on the Superion device in its recommendation.”

The overall impressions summarized by the authors identified suggested improvements at up to 3 years follow-up (based on four studies); however, current evidence does not inform whether clinical outcomes associated with the Superion Interspinous Spacer are the same, better, or worse than other currently available minimally invasive interventions, or with more established surgeries involving spinal fusion. Although one randomized controlled trial was identified, it compares Superion with a product no longer on the U.S. market (X-Stop). Two registered ongoing studies do not have control groups and are therefore unlikely to address the gaps in this evidence base. No clinical practice guidelines address Superion specifically, and guidance on use of spacers in general is mixed.

Additionally, final outcomes from a single RCT with the Superion device (FDA noninferiority trial versus X-stop) was reported in four publications and post-hoc analysis in three additional reports.²⁷⁻³³ This single-study with five-year follow-up initially randomized 189 patients to the Superion treatment group. Interpretations from these study data are limited by the lack of blinding and lack of control groups treated by surgical decompression or medical management. Additionally, the FDA-mandated primary endpoint of this trial was non-inferiority to X-STOP at 2 years, with additional postmarket surveillance for 10 years. As stated in the Hayes report, the comparison group (X-Stop) has been pulled off the U.S. market.

CLINICAL PRACTICE GUIDELINES

Dynamic Stabilization Devices

North American Spine Society (NASS)

In 2016, the NASS addressed dynamic stabilization without arthrodesis for the treatment of degenerative lumbar spondylolisthesis.^{34,35} The workgroup made no recommendation due to the lack of quality evidence. Authors called for large, prospective studies with long-term follow-up comparing dynamic stabilization to medical or interventional treatment.

Interspinous Spacers

North American Spine Society (NASS)

In 2025, the North American Spine Society (NASS) published a coverage policy recommendation for lumbar interspinous device without fusion and with decompression.³⁶

“There are several interspinous distraction devices available on the market today. These include static (ie, nonflexible or noncompressible) devices and dynamic implants. Static devices are used to provide indirect decompression of the neural elements. Dynamic devices, according to their FDA labeling, are intended to be used in conjunction with a direct decompression (eg, laminectomy). For the purpose of this document, coverage recommendations will apply only to static (ie, nonflexible or noncompressible) devices that are intended to be used instead of a direct decompressive procedure.

“The initially introduced interspinous devices (ISDs) have lost popularity for a variety of reasons; newer devices seemed to have filled their place. Since the publication of the original coverage document, there have been a number of additional studies performed, some of which supersede prior data. Based on an updated review of the current evidence-base, interspinous distraction devices without fusion may be indicated for the following diagnoses with qualifying criteria, when appropriate:

“Clinically symptomatic degenerative lumbar stenosis:

- associated with neurogenic claudication that is relieved by lumbar flexion
- sitting for at least 50 minutes without pain
- can walk at least 50 feet
- mild to moderate stenosis as assessed on advanced imaging (CT or MRI) defined as 25-50% reduction in lateral/ central foraminal diameter compared to adjacent levels
- failure of 6 months nonoperative treatment
- no more than 25° of degenerative scoliosis
- no more than a Grade I degenerative spondylolisthesis
- open surgery (eg, laminectomy) is not a medically safe treatment option because of comorbidities

“Interspinous distraction devices are NOT indicated in cases that do not fall within the above parameters. In particular, they are not indicated in the following scenarios and conditions:

- degenerative spondylolisthesis of Grade II or higher
- degenerative scoliosis greater than 25°
- dynamic instability at the operative level
- symptoms are not relieved by flexion”

In 2014, NASS published revised clinical guidelines for degenerative lumbar spondylolisthesis.³⁵ For the clinical question, “Does the use of interspinous spacers in the treatment of degenerative lumbar spondylolisthesis improve outcomes compared to nonoperative treatment,” the guidelines concluded: There is insufficient and conflicting evidence to make a recommendation for or against the efficacy of interspinous spacers versus medical/ interventional treatment in the management of degenerative lumbar spondylolisthesis patients. (Grade of Recommendation I - Insufficient Evidence). The authors stated a need for large studies with long-term follow-up comparing interspinous spacers to medical/interventional treatment to better assess the safety and efficacy of interspinous spacers.

The 2011 revised clinical guidelines from NASS on lumbar spinal stenosis stated: There is insufficient evidence at this time to make a recommendation for or against the placement of an interspinous process spacing device in patients with lumbar spinal stenosis.³⁷ (Grade of Recommendation I - Insufficient Evidence)

EVIDENCE SUMMARY

Evidence does not support the safety and efficacy of dynamic stabilization devices and interspinous spacers. All systematic reviews to date note a paucity of long-term evidence from high-quality trials. The literature largely comprises small, uncontrolled studies with short-term follow-up. Moreover, no evidence-based clinical practice guidelines recommend dynamic stabilization devices. While one guideline recommends interspinous spacers in addition to interbody fusion and/or decompression fusion for the treatment of lumbar spinal stenosis, the evidence used to support this recommendation

suffers from the following limitations: heterogenous patient populations; nonrandomized design; high potential for patient selection bias; relatively limited follow-up in most studies; small sample size; absence of power analyses; inadequate blinding; potential for reporting bias, including censored data and missing data for some outcomes and time point.

HEALTH EQUITY CONSIDERATIONS

The Centers for Disease Control and Prevention (CDC) defines health equity as the state in which everyone has a fair and just opportunity to attain their highest level of health. Achieving health equity requires addressing health disparities and social determinants of health. A health disparity is the occurrence of diseases at greater levels among certain population groups more than among others. Health disparities are linked to social determinants of health which are non-medical factors that influence health outcomes such as the conditions in which people are born, grow, work, live, age, and the wider set of forces and systems shaping the conditions of daily life. Social determinants of health include unequal access to health care, lack of education, poverty, stigma, and racism.

The U.S. Department of Health and Human Services Office of Minority Health calls out unique areas where health disparities are noted based on race and ethnicity. Providence Health Plan (PHP) regularly reviews these areas of opportunity to see if any changes can be made to our medical or pharmacy policies to support our members obtaining their highest level of health. Upon review, PHP creates a Coverage Recommendation (CORE) form detailing which groups are impacted by the disparity, the research surrounding the disparity, and recommendations from professional organizations. PHP Health Equity COREs are updated regularly and can be found online [here](#).

BILLING GUIDELINES AND CODING

The following codes do not apply to minimally invasive dynamic stabilization procedures of the spine and should not be used to bill for these services: 22533, 22534, 22558, 22585, 22586, 22612, 22614, 22630, 22632, 22633, 22634, 22800, 22802, 22804, 22808, 22810, 22812, 22840, 22841, 22842, 22843, 22844, 22845, 22846, 22847, 22853, 22854, 22859.

CODES*		
CPT	22867	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level
	22868	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (List separately in addition to code for primary procedure)
	22869	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level

	22870	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (List separately in addition to code for primary procedure)
	22899	Unlisted procedure, spine
	64999	Unlisted procedure, nervous system
HCPCS	C1821	Interspinous process distraction device (implantable)
	L8699	Prosthetic implant, not otherwise specified

***Coding Notes:**

- The above code list is provided as a courtesy and may not be all-inclusive. Inclusion or omission of a code from this policy neither implies nor guarantees reimbursement or coverage. Some codes may not require routine review for medical necessity, but they are subject to provider contracts, as well as member benefits, eligibility and potential utilization audit.
- All unlisted codes are reviewed for medical necessity, correct coding, and pricing at the claim level. If an unlisted code is submitted for non-covered services addressed in this policy then it will be **denied as not covered**. If an unlisted code is submitted for potentially covered services addressed in this policy, to avoid post-service denial, **prior authorization is recommended**.
- See the non-covered and prior authorization lists on the Company [Medical Policy, Reimbursement Policy, Pharmacy Policy and Provider Information website](#) for additional information.
- HCPCS/CPT code(s) may be subject to National Correct Coding Initiative (NCCI) procedure-to-procedure (PTP) bundling edits and daily maximum edits known as “medically unlikely edits” (MUEs) published by the Centers for Medicare and Medicaid Services (CMS). This policy does not take precedence over NCCI edits or MUEs. Please refer to the CMS website for coding guidelines and applicable code combinations.

REFERENCES

1. Zimmer Biomet, Aspen® MIS Fusion System. https://thespinemarketgroup.com/wp-content/uploads/2021/03/1079.2-US-en-REV1020_Aspen-Surgical-Technique.pdf. Accessed 4/20/2025.
2. Surgalign, DSS® Stabilization System. https://www.surgalign.com/en_gb/product/dss-dynamic-stabilization-system/. Accessed 4/20/2025.
3. Zimmer Biomet, Dynesys Top-Loading System. <https://thespinemarketgroup.com/wp-content/uploads/2015/10/dynesys-top-loading-system-surgical-technique.pdf>. Accessed 4/20/2025.
4. Surgalign, Coflex. <https://coflexsolution.com/>. Accessed 4/20/2025.
5. Food and Drug Administration. Superion Interspinous Spacer. Decision Date, 5/20/2015. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P140004>. Accessed 4/20/2025.
6. Boston Scientific | The Vertiflex™ Procedure | Superion™ Indirect Decompression System. <https://www.bostonscientific.com/en-US/products/indirect-decompression-system/superion-indirect-decompression-system.html>. Accessed 4/20/2025.
7. FDA. Premarket Approval (PMA). Device: X Stop Interspinous Process Decompression System Withdrawl date: 04/30/2015. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P040001>. Accessed 4/20/2025.
8. Food and Drug Administration. Biomet Fusion System. Zimmer Biomet Spine Inc. Decision Date 2/28/2017. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P040001>. Accessed 4/20/2025.

9. Food and Drug Administration. DSS Stabilization System. Decision Date 7/2/2010. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K101083>. Accessed 4/20/2025.
10. Food and Drug Administration. Dynesys Spinal System. Decision Date, 3/5/2004. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K031511>. Accessed 4/20/2025.
11. Food and Drug Administration. Isobar Spinal System. Decision Date, 1/15/2002. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K013447>. Accessed 4/20/2025.
12. Food and Drug Administration. Coflex Interlaminar Technology. Decision Date, 2/13/2012. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmn.cfm?ID=K112595>. Accessed 4/20/2025.
13. Kaibara T, Karahalios DG, Porter RW, et al. Biomechanics of a lumbar interspinous anchor with transforaminal lumbar interbody fixation. *World neurosurgery*. 2010;73(5):572-577.
14. Karahalios DG, Kaibara T, Porter RW, et al. Biomechanics of a lumbar interspinous anchor with anterior lumbar interbody fusion. *Journal of Neurosurgery: Spine*. 2010;12(4):372-380.
15. Bieri KS, Goodwin K, Aghayev E, Riesner H-J, Greiner-Perth R. Dynamic Posterior Stabilization versus Posterior Lumbar Intervertebral Fusion: A Matched Cohort Study Based on the Spine Tango Registry. *Journal of Neurological Surgery Part A: Central European Neurosurgery*. 2018;79(03):224-230.
16. Lee C-H, Jahng T-A, Hyun S-J, et al. Dynamic stabilization using the Dynesys system versus posterior lumbar interbody fusion for the treatment of degenerative lumbar spinal disease: a clinical and radiological outcomes-based meta-analysis. *Neurosurgical focus*. 2016;40(1):E7.
17. Qian J, Bao Z, Li X, Zou J, Yang H. Short-term therapeutic efficacy of the ISOBAR TTL dynamic internal fixation system for the treatment of lumbar degenerative disc diseases. *Pain physician*. 2016;19(6):E853-861.
18. Xing R, Dou Q, Li X, et al. Posterior dynamic stabilization with direct pars repair via Wiltse approach for the treatment of lumbar spondylolysis: the application of a novel surgery. *Spine*. 2016;41(8):E494-E502.
19. Fu L, France A, Xie Y, Fang K, Gan Y, Zhang P. Functional and radiological outcomes of semi-rigid dynamic lumbar stabilization adjacent to single-level fusion after 2 years. *Archives of orthopaedic and trauma surgery*. 2014;134(5):605-610.
20. Hayes Inc. Coflex Interlaminar Stabilization Device (Surgalign Spine Technologies Inc.) for Treatment of Lumbar Spinal Stenosis. <https://evidence.hayesinc.com/report/htb.coflex2708>. Published 2024. Accessed 4/20/2025.
21. Mo Z, Li D, Zhang R, Chang M, Yang B, Tang S. Comparative effectiveness and safety of posterior lumbar interbody fusion, Coflex, Wallis, and X-stop for lumbar degenerative diseases: A systematic review and network meta-analysis. *Clinical neurology and neurosurgery*. 2018.
22. Zhao X-w, Ma J-x, Ma X-l, et al. Interspinous process devices (IPD) alone versus decompression surgery for lumbar spinal stenosis (LSS): A systematic review and meta-analysis of randomized controlled trials. *International Journal of Surgery*. 2017;39:57-64.
23. Machado GC, Ferreira PH, Yoo RI, et al. Surgical options for lumbar spinal stenosis. *Cochrane Database of Systematic Reviews*. 2016(11).
24. Moojen WA, Arts MP, Jacobs WC, et al. IPD without bony decompression versus conventional surgical decompression for lumbar spinal stenosis: 2-year results of a double-blind randomized controlled trial. *European Spine Journal*. 2015;24(10):2295-2305.

25. ECRI. Superion Indirect Decompression System (Boston Scientific Corp.) for Treating Lumbar Spinal Stenosis. Updated 8/19/2022. <https://members.ecri.org/evidenceanalysis/superion-indirect-decompression-system-boston-scientific-corp-for-treating->. Accessed 4/20/2025.
26. Hayes Inc. Superion Interspinous Spacer System (Vertiflex) for Treatment of Neurogenic Claudication Caused by Spinal Stenosis. <https://evidence.hayesinc.com/report/eer.superion4813>. Published 2024. Accessed 4/20/2025.
27. Nunley PD, Deer TR, Benyamin RM, Staats PS, Block JE. Interspinous process decompression is associated with a reduction in opioid analgesia in patients with lumbar spinal stenosis. *J Pain Res*. 2018;11:2943-2948. <https://www.ncbi.nlm.nih.gov/pubmed/30538533>
28. Nunley PD, Patel VV, Orndorff DG, Lavelle WF, Block JE, Geisler FH. Interspinous Process Decompression Improves Quality of Life in Patients with Lumbar Spinal Stenosis. *Minim Invasive Surg*. 2018;2018:1035954. <https://www.ncbi.nlm.nih.gov/pubmed/30057811>
29. Nunley PD, Patel VV, Orndorff DG, Lavelle WF, Block JE, Geisler FH. Five-year durability of stand-alone interspinous process decompression for lumbar spinal stenosis. *Clin Interv Aging*. 2017;12:1409-1417. <https://www.ncbi.nlm.nih.gov/pubmed/28919727>
30. Patel VV, Nunley PD, Whang PG, et al. Superion((R)) InterSpinous Spacer for treatment of moderate degenerative lumbar spinal stenosis: durable three-year results of a randomized controlled trial. *J Pain Res*. 2015;8:657-662. <https://www.ncbi.nlm.nih.gov/pubmed/26491369>
31. Patel VV, Whang PG, Haley TR, et al. Superion interspinous process spacer for intermittent neurogenic claudication secondary to moderate lumbar spinal stenosis: two-year results from a randomized controlled FDA-IDE pivotal trial. *Spine (Phila Pa 1976)*. 2015;40(5):275-282. <https://www.ncbi.nlm.nih.gov/pubmed/25494323>
32. Patel VV, Whang PG, Haley TR, et al. Two-year clinical outcomes of a multicenter randomized controlled trial comparing two interspinous spacers for treatment of moderate lumbar spinal stenosis. *BMC Musculoskelet Disord*. 2014;15:221. <https://www.ncbi.nlm.nih.gov/pubmed/24996648>
33. Miller LE, Block JE. Interspinous spacer implant in patients with lumbar spinal stenosis: preliminary results of a multicenter, randomized, controlled trial. *Pain Res Treat*. 2012;2012:823509. <https://www.ncbi.nlm.nih.gov/pubmed/22448323>
34. Matz PG, Meagher RJ, Lamer T, et al. Guideline summary review: An evidence-based clinical guideline for the diagnosis and treatment of degenerative lumbar spondylolisthesis. *Spine J*. 2016;16(3):439-448.
35. North American Spine Society. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care. Diagnosis and Treatment of Degenerative Lumbar Spondylolisthesis. <https://www.spine.org/Documents/ResearchClinicalCare/Guidelines/Spondylolisthesis.pdf>. Published 2014. Accessed 4/20/2025.
36. North American Spine Society. Interspinous Device without Fusion and with Decompression. <https://www.spine.org/coverage>. Published 2025. Accessed 4/20/2025.
37. North American Spine Society. Evidence-Based Clinical Guidelines for Multidisciplinary Spine Care. Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LumbarStenosis.pdf>. Published 2011. Accessed 4/20/2025.

POLICY REVISION HISTORY

DATE	REVISION SUMMARY
2/2023	Converted to new policy template.

6/2023	Change denial from investigational to not medically necessary.
7/2023	Annual update, no changes.
6/2024	Annual update. Policy title change. No changes to coding or criteria.
9/2024	Interim update. Correction of codes listed for dynamic stabilization devices and interspinous spacers.
6/2025	Annual update. No changes to criteria.