


MEDICAL POLICY	Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)
Effective Date: 08/01/2022	Medical Policy Number: 204
 8/1/2022	Technology Assessment Committee Approved Date: 6/16 Medical Policy Committee Approved Date: 8/92; 4/95; 8/97; 1/98; 1/00; 1/01; 2/03; 10/03; 9/04; 11/05; 11/07; 11/08; 7/10; 2/11; 6/13; 4/14; 2/15; 8/15; 8/17; 5/18; 3/19; 5/19; 06/2020; 07/2021; 7/2022
Medical Officer	Date

See Policy CPT/HCPCS CODE section below for any prior authorization requirements

SCOPE:

Providence Health Plan, Providence Health Assurance, Providence Plan Partners, and Ayn Health Solutions as applicable (referred to individually as “Company” and collectively as “Companies”).

APPLIES TO:

All lines of business except Medicare

BENEFIT APPLICATION

Medicaid Members

Oregon: Services requested for Oregon Health Plan (OHP) members follow the OHP Prioritized List and Oregon Administrative Rules (OARs) as the primary resource for coverage determinations. Medical policy criteria below may be applied when there are no criteria available in the OARs and the OHP Prioritized List.

POLICY CRITERIA

I. The following criteria are based on the Undersea and Hyperbaric Medical Society’s (UHMS) Hyperbaric Oxygen Therapy Indications: 14th Edition.^{1,2} Hyperbaric oxygen therapy (HBOT), performed within the UHMS treatment guidelines, may be considered **medically necessary** for the following UHMS indications:

UHMS Indication	UHMS Treatment Guideline
Acute thermal burn injury	30 sessions; UHMS indicates it is rare to exceed 40-50 sessions
Air or gas embolism	2 sessions; UHMS indicates 5-10 sessions may be necessary

MEDICAL POLICY

Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)

<p>Arterial insufficiencies (At least one of the following criteria [A. <u>or</u> B] must be met):</p> <p>A. Diabetic lower extremity wounds when all of the following (1.-4.) additional criteria are met:</p> <ol style="list-style-type: none"> 1. Patient with Type 1 or Type 2 Diabetes with lower extremity wound due to diabetes; and 2. Wagner grade III or higher wound severity (see Policy Guidelines section); and 3. Patient has failed adequate course of standard wound therapy (see Policy Guidelines section); and 4. Re-evaluations at 30 days must show continued progress; or <p>B. Arterial insufficiency ulcer when at least one of the following (1.-2.) criteria are met:</p> <ol style="list-style-type: none"> 1. The patient has persistent hypoxia despite attempts at increasing blood flow; or 2. Wound failure continues despite maximum revascularization. 	<p>Treatment varies depending upon the severity of the wound and the type of chamber used.</p> <ul style="list-style-type: none"> - For multiplace and monoplace chambers UHMS recommends 90-120 minute sessions once or twice daily (i.e., 30-60 sessions in a 30 day time span). - When stabilized, once daily treatment is recommended.
<p>Carbon monoxide poisoning</p>	<p>5 sessions</p>
<p>Central Retinal Artery Occlusion (CRAO) when HBOT treatment is initiated within 24 hours of vision loss</p>	<ul style="list-style-type: none"> - If vision shows improvement, treat with 90 minutes sessions for a minimum of 3 days. - Continue treatment until there is three consecutive days with no clinical improvement.
<p>Compartment syndrome</p>	<ul style="list-style-type: none"> - Twice a day for 24-36 hours with oxygen breathing for 90 minutes each, or a single treatment a day for 120 minutes. - For residual complications after fasciotomy, treatments should be twice a day for 7-10 days, or when condition is stabilized such that no additional benefit is received.
<p>Compromised skin grafts and flaps</p>	<ul style="list-style-type: none"> - Initial treatment is for 90-120 minutes. - Once the flap or graft is stable, once daily treatments may suffice.
<p>Crush injuries</p>	<ul style="list-style-type: none"> - Two or more treatments a day with oxygen breathing for 90 minutes each, or a single treatment a day for 120 minutes.

MEDICAL POLICY

Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)

Cyanide poisoning	5 sessions
Decompression sickness	10 sessions
Delayed radiation injury (soft tissue and bony necrosis)	30 sessions
Gas gangrene (Clostridial myositis and myonecrosis)	10 sessions
Idiopathic sudden sensorineural hearing loss: moderate to profound (≥ 41 dB) when HBOT treatment is initiated within 14 days of symptom onset.	20 sessions
Intracranial abscess (includes cerebral abscess, subdural empyema, and epidural empyema) when HBOT is used as an adjunctive therapy in patients who meet at least one of the following (A.-E.) criteria: A. Multiple abscesses; or B. Abscesses in a deep or dominant location; or C. Immune compromised; or D. In situations where surgery is contraindicated or where the patient is a poor surgical risk; or E. No response or further deterioration in spite of standard surgical (e.g., 1-2 needle aspirates) and antibiotic treatment.	– Treatment should be administered for 60-90 minutes once or twice daily, depending upon the severity of the condition.
Necrotizing soft-tissue infections	– Treatment is given for 90 minutes twice daily during the initial phase of therapy, until there is no longer evidence of progression and infection is under control. – Once the patient’s condition is stabilized, and prior to treatment cessation, treatment once daily may be instituted to assure relapse will not occur.
Radiation necrosis	60 sessions
Refractory osteomyelitis (stage 3B and 4B)(see Policy Guidelines section)	40 postoperative sessions over a 4-6 week period
Severe anemia when transfusion is not possible	HBOT therapy should be continued with taper of both time and frequency until red blood cells have been replaced by patient regeneration or the patient can undergo blood transfusion.

II. Hyperbaric oxygen therapy is considered **investigational and is not covered** when criterion I. above is not met, including, but not limited to any of the following:

- A. Acute coronary syndrome
- B. Acute ischemic stroke
- C. Acute surgical and traumatic wounds

MEDICAL POLICY

Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)

- D. AIDS/HIV
- E. Alzheimer's disease
- F. Asthma
- G. Autism Spectrum Disorder
- H. Bell's Palsy
- I. Blindness
- J. Brain injury including traumatic (TBI) and chronic brain injury
- K. Cerebral Palsy
- L. Concurrent treatment with other non-standard wound care (e.g., wound vac, negative pressure wound therapy)
- M. Delayed onset muscle soreness and closed soft tissue injury
- N. Depression
- O. Fracture healing
- P. Headache- migraine and cluster
- Q. Heart disease
- R. Hepatitis
- S. Lower extremity injury (e.g., sprain, tendonitis, fracture, dislocation)
- T. Multiple Sclerosis
- U. Non-healing lower extremity wound (e.g., ischemic ulcer) with no arterial blood flow
- V. Otitis externa
- W. Parkinson's disease
- X. Perianal fistulas
- Y. Posttraumatic stress disorder and acute stress disorder
- Z. Pressure ulcers
- AA. Shoulder injury
- BB. Spinal cord injury
- CC. Tumor sensitization to radiotherapy
- DD. Vascular dementia
- EE. Venous ulcers
- FF. Wound caused by or not healing due to a foreign body reaction (e.g., mesh, suture)

III. Topical hyperbaric oxygen therapy is considered **not medically necessary and is not covered** for all indications.

POLICY GUIDELINES

Wagner Grading System for Diabetic Foot Infections³

- Grade 0 - Intact Skin
- Grade 1 - Superficial ulcer of skin or subcutaneous tissue
- Grade 2 - Ulcers extend into tendon, bone, or capsule
- Grade 3 - Deep ulcer with osteomyelitis, or abscess
- Grade 4 - Gangrene of toes or forefoot

- Grade 5 - Midfoot or hindfoot gangrene

Standard Wound Therapy¹

Defined as 30 days of treatment including assessment and correction of vascular abnormalities, optimization of nutritional status and glucose control, debridement, moist wound dressing, off-loading, and treatment of infection.

Osteomyelitis Staging⁴

Anatomic type

- **Stage 1:** Medullary osteomyelitis

Medullary osteomyelitis denotes infection confined to the intramedullary surfaces of the bone. Hematogenous osteomyelitis and infected intramedullary rods are examples of this anatomic type.

- **Stage 2:** Superficial osteomyelitis

Superficial osteomyelitis is a true contiguous focus infection of bone; it occurs when an exposed infected necrotic surface of bone lies at the base of a soft-tissue wound.

- **Stage 3:** Localized osteomyelitis

Localized osteomyelitis is usually characterized by a full thickness, cortical sequestration which can be removed surgically without compromising bony stability.

- **Stage 4:** Diffuse osteomyelitis

Diffuse osteomyelitis is a through-and-through process that usually requires an intercalary resection of the bone to arrest the disease process. Diffuse osteomyelitis includes those infections with a loss of bony stability either before or after debridement surgery.

Physiologic class of host

- **Class A** denotes a normal host
- **Class B** denotes a host with systemic compromise, local compromise, or both
- **Class C** denotes a host for whom the morbidity of treatment is worse than that imposed by the disease itself

MEDICAL POLICY	Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)
-----------------------	--

CPT/HCPCS CODES

All Lines of Business Except Medicare	
Prior Authorization Required	
99183	Physician or other qualified health care professional attendance and supervision of hyperbaric oxygen therapy, per session
G0277	Hyperbaric oxygen under pressure, full body chamber, per 30 minute interval
Not Covered	
A4575	Topical hyperbaric oxygen chamber, disposable
E0446	Topical oxygen delivery system, not otherwise specified, includes all supplies and accessories

DESCRIPTION

Hyperbaric Oxygen Therapy (HBOT)

The Undersea and Hyperbaric Medical Society (UHMS) defines hyperbaric oxygen therapy (HBOT) as “an intervention in which an individual breathes near 100% oxygen intermittently while inside a hyperbaric chamber that is pressurized to greater than sea level pressure.”² For certain indications, HBOT is the primary treatment modality while in other indications it is an adjunctive treatment to surgical or pharmacological interventions. Clinical treatments may take place in a Class A (multi-chamber) or Class B (mono-chamber) system. A Class A system holds two or more people while a Class B system holds only the patient.

UHMS Approved HBOT Indications

UHMS Indication	Description
Acute thermal burn injury	Anatomic, physiologic, endocrinologic, and immunologic alterations due to a burn injury. ⁵
Air or gas embolism	When one or more air bubbles enter a vein or artery and block it. ⁶
Diabetic lower extremity wounds	An open sore or ulcer, most commonly located on the bottom of the foot, caused by diabetes-related circulatory issues. ⁷
Carbon monoxide poisoning	Carbon monoxide is an odorless, colorless gas that can be deadly upon exposure. ⁸
Central Retinal Artery Occlusion (CRAO)	A disease of the eye where the flow of blood through the central retinal artery is blocked (occluded). ⁹
Compartment syndrome	A condition that occurs when pressure within the muscles builds to dangerous levels. This pressure can decrease blood flow, which prevents nourishment and oxygen from reaching nerve and muscle cells. ¹⁰
Compromised skin grafts and flaps	Skin grafts and flaps is a technique used in reconstructive surgery where a type of tissue is lifted from a donor site and moved to a

MEDICAL POLICY	Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)
-----------------------	--

	recipient site. In tissue compromised by irradiation or decreased oxygen supply, HBOT is used to maximize viability of the graft or flap. ¹¹
Crush injuries	An injury that occurs when force or pressure is put on a body part. This injury happens when part of the body is compressed between two heavy objects. ¹²
Cyanide poisoning	Cyanide toxicity is a rare form of poisoning due to exposure to cyanide. Cyanide exposure occurs relatively frequently in patients with smoke inhalation. ¹³
Decompression sickness	Injuries caused by a rapid increase in the pressure that surrounds you, of either air or water. It occurs most commonly in scuba or deep-sea divers, although it also can occur during high-altitude or unpressurized air travel. ¹⁴
Gas gangrene	A highly lethal soft tissue infection of skeletal muscle caused by toxin and gas producing <i>Clostridium</i> bacteria species. ¹⁵
Intracranial abscess (includes cerebral abscess, subdural empyema, and epidural empyema)	A collection of pus, immune cells, and other material in the brain, usually from a bacterial or fungal infection. ¹⁶
Necrotizing soft-tissue infections	A rare but severe type of bacterial infection that can destroy the muscles, skin, and underlying tissue. ¹⁷
Radiation necrosis	Damage done to non-osseous tissues by ionizing radiation during the course of radiotherapy for cancer. ¹⁸
Refractory osteomyelitis (stage 3B and 4B)	A bone infection that has not responded to appropriate medical treatment (refractory). ¹⁹
Severe anemia when transfusion is not possible	Hemoglobin concentrations below 8.0 g/dL. ²⁰ Some religions prevent people from receiving blood transfusions.
Idiopathic sudden sensorineural hearing loss (ISSHL)	Unexplained unilateral hearing loss with onset over a period of less than 72 hours. ²¹

Topical Oxygen Therapy (TOT)

TOT is intended to increase wound oxygenation and promote wound healing. There are two types of TOT:

- Hyperbaric TOT (HTOT): “The affected limb is enclosed in a chamber or gas-impermeable bag, and the chamber is filled with oxygen pressurized slightly above atmospheric pressure. HTOT requires patient immobility during in-clinic treatment sessions, which may last 90 minutes once per day for weeks.”²²
- Continuous TOT (CTOT): An alternative to HTOT that does not require patient immobilization or in-clinic administration. CTOT can also be used at the same time as dressings and offloading. “A portable oxygen concentrator refines and delivers atmospheric (normobaric) oxygen to the wound site through a cannula.”²²

REVIEW OF EVIDENCE

The medically necessary indications for hyperbaric oxygen therapy is based on the Undersea and Hyperbaric Medical Society's (UHMS) Hyperbaric Oxygen Therapy Indications: 13th Edition.^{1,2} Therefore, an evidence review was not conducted for these indications.

A review of the ECRI, Hayes, Cochrane, and PubMed databases was conducted regarding the use of hyperbaric oxygen therapy for indications not included in the UHMS recommendation. A review of evidence was also conducted for the use of topical hyperbaric oxygen therapy. Below is a summary of the available evidence identified through June 2022.

Investigational Indications for Hyperbaric Oxygen Therapy

All of the following purported indications for hyperbaric oxygen therapy (HBOT) are considered investigational. This investigational stance is supported by one, or more, of the following (1) a Cochrane systematic review (2) an evidence-based clinical practice guideline, and/or (3) an FDA consumer warning against the use of HBOT for that indication. See the associated reference for more information.

- Acute coronary syndrome^{23,24}
- Acute ischemic stroke^{25,26}
- Acute surgical and traumatic wounds²⁷
- AIDS/HIV²⁵
- Alzheimer's disease²⁵
- Asthma²⁵
- Autism Spectrum Disorder²⁸⁻³⁰
- Bell's Palsy^{24,25,31}
- Brain injury including traumatic (TBI) and chronic brain injury^{24,25,32}
- Cerebral Palsy²⁵
- Delayed onset muscle soreness and closed soft tissue injury³³
- Depression²⁵
- Fracture healing^{24,34}
- Headache- migraine and cluster^{24,25,35}
- Heart disease²⁵
- Hepatitis²⁵
- Lower extremity injury (e.g., sprain, tendonitis, fracture, dislocation)³⁶
- Multiple Sclerosis^{25,37}
- Otitis externa^{24,38}
- Parkinson's disease²⁵
- Posttraumatic stress disorder and acute stress disorder³⁹
- Pressure ulcers⁴⁰
- Shoulder injury⁴¹
- Spinal cord injury²⁵
- Tumor sensitization to radiotherapy⁴²
- Vascular dementia^{24,43}

- Venous ulcers²⁴

Topical Hyperbaric Oxygen Therapy

As indicated by the evidence review, topical hyperbaric oxygen therapy (HBOT) has been predominantly investigated as a treatment of chronic wounds.⁴⁴⁻⁵³ In 2022, Hayes updated an evidence review evaluating topical oxygen therapy for chronic wound healing.²² The review identified three randomized controlled trials (RCT) as eligible for inclusion. Sample sizes ranged from 20 to 130 patients and follow-up times varied from 8 to 12 weeks. Outcomes of interest included complete wound healing, time to complete wound healing, and complications.

The results indicated that topical hyperbaric oxygen therapy may provide an incremental benefit to standard wound care for healing chronic diabetic foot ulcers that have failed to respond to wound care alone. However, not all studies reported a benefit and there is still insufficient outcome and safety data to inform meaningful conclusions. Additionally, all studies evaluated topical HBOT for chronic diabetic foot ulcers; therefore, the evidence is insufficient to inform evidence-based conclusions regarding topical HBOT for other types of chronic wounds.

Hayes determined the available evidence to be of low quality. Hayes concluded the following ratings:

- D2 (insufficient evidence)—For the use of continuous topical oxygen therapy in adults with diabetes-related foot ulcers that are refractory to standard wound care.
- D2 (insufficient evidence)—For the use of continuous topical oxygen therapy for any other chronic wound type other than DFU.
- D2 (insufficient evidence)—For the use of hyperbaric topical oxygen therapy for any chronic wound type.²²

CLINICAL PRACTICE GUIDELINES

Undersea and Hyperbaric Medical Society (UHMS)

The UHMS 2019 Hyperbaric Oxygen Therapy Indications recommends the following (A.-Q.) indications and treatment guidelines for systemic hyperbaric oxygen therapy (HBOT).^{1,2}

- Air or gas embolism
- Carbon monoxide poisoning
- Carbon monoxide poisoning complicated by cyanide poisoning
- Central retinal artery occlusion
- Clostridial myositis and myonecrosis (gas gangrene)
- Crush injury, compartment syndrome, and other acute traumatic ischemias
- Decompression sickness
- Enhancement of healing in select problem wounds
- Exceptional blood loss (severe anemia)
- Intracranial abscess
- Necrotizing soft tissue infections
- Osteomyelitis (refractory)

MEDICAL POLICY	Hyperbaric Oxygen Therapy (All Lines of Business Except Medicare)
-----------------------	--

- Delayed radiation injury (soft tissue and bony necrosis)
- Skin grafts and flaps (compromised)
- Thermal burns (acute)
- Idiopathic sudden sensorineural hearing loss

National Institute for Health and Care Excellence (NICE)

In 2014, the NICE published a guideline on prevention and management of pressure ulcers.⁵⁴ As part of these recommendations, investigators recommended that HBO not be used as a treatment modality for pressure ulcers in adults, children, infants or neonates. No randomized controlled trials were identified evaluating HBO for pressure ulcers.

POLICY SUMMARY

The medically necessary indications for hyperbaric oxygen therapy are based on the Undersea and Hyperbaric Medical Society's (UHMS) Hyperbaric Oxygen Therapy Indications: 14th Edition. Additionally, the FDA follows UHMS for determining the FDA-approved indications for HBOT. There are numerous investigational indications for hyperbaric oxygen therapy. There is insufficient published evidence to adequately evaluate the efficacy and/or safety of these indications. Additional good-quality research, as well as approval by the UHMS and FDA, is required to support other purported indications for HBOT. There is also insufficient evidence to permit reliable conclusions regarding topical HBOT. Further studies of good methodological quality are required to establish the efficacy and safety of topical HBOT. Additional studies also need to demonstrate an improvement in patient health outcomes with topical HBOT compared to standard, systemic HBOT.

INSTRUCTIONS FOR USE

Company Medical Policies serve as guidance for the administration of plan benefits. Medical policies do not constitute medical advice nor a guarantee of coverage. Company Medical Policies are reviewed annually and are based upon published, peer-reviewed scientific evidence and evidence-based clinical practice guidelines that are available as of the last policy update. The Companies reserve the right to determine the application of Medical Policies and make revisions to Medical Policies at any time. Providers will be given at least 60-days notice of policy changes that are restrictive in nature.

The scope and availability of all plan benefits are determined in accordance with the applicable coverage agreement. Any conflict or variance between the terms of the coverage agreement and Company Medical Policy will be resolved in favor of the coverage agreement.

REGULATORY STATUS

U.S. Food and Drug Administration (FDA)

The FDA approved indications for hyperbaric oxygen therapy (HBOT) are based on the Undersea and Hyperbaric Medical Society (UHMS) recommended indications for HBOT.⁵⁵ Therefore, HBOT is FDA-approved for the following indications:

- Acute Thermal Burn Injury
- Air or Gas Embolism
- Arterial Insufficiencies
- Carbon Monoxide Poisoning
- Central Retinal Artery Occlusion (CRAO)
- Compartment syndrome
- Compromised Skin Grafts and Flaps
- Crush Injuries
- Cyanide Poisoning Decompression Sickness
- Gas Gangrene
- Intracranial Abscess (includes cerebral abscess, subdural empyema, and epidural empyema)
- Necrotizing Soft-Tissue Infections
- Radiation Necrosis
- Refractory Osteomyelitis
- Severe Anemia
- Idiopathic Sudden Sensorineural Hearing Loss (ISSHL)

Mental Health Parity Statement

Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case. In cases where medical necessity is not established by policy for specific treatment modalities, evidence not previously considered regarding the efficacy of the modality that is presented shall be given consideration to determine if the policy represents current standards of care.

REFERENCES

1. Undersea and Hyperbaric Medical Society. *Hyperbaric Oxygen Therapy Indications: The Hyperbaric Oxygen Therapy Committee Report*. Best Publishing Company; 2019.
2. Richard E. Moon. Undersea and Hyperbaric Oxygen Medical Society: Hyperbaric Oxygen Therapy Indications 14th Edition - Background and References. <https://www.uhms.org/images/UHMS-Reference-Material.pdf>. Published 2019. Accessed 6/27/2022.
3. Wheelless' Textbook of Orthopedics. http://www.wheellesonline.com/ortho/wagner_grading_system_for_diabetic_foot_infections. Accessed 6/27/2022.
4. The Osteomyelitis Center of Central Florida: Staging System for Long Bone Osteomyelitis. <https://www.osteomyelitiscenter.com/staging-of-osteomyelitis.html>. Accessed 6/27/2022.

5. Garner WL, Magee W. Acute burn injury. *Clinics in plastic surgery*. 2005;32(2):187-193.
6. Healthline: What is an air embolism? <https://www.healthline.com/health/air-embolism>. Published 2018. Accessed 6/27/2022.
7. Memorial Herman Wound Care: Diabetic Wounds of the Lower Extremities. <http://www.memorialhermann.org/wound-care/diabetic-foot-wounds/>. Accessed 6/27/2022.
8. The Centers for Disease Control and Prevention. Carbon Monoxide Poisoning. <https://www.cdc.gov/co/faqs.htm>. Accessed 6/27/2022.
9. Medscape. Central Retinal Artery Occlusion. <https://emedicine.medscape.com/article/1223625-overview>. Accessed 6/17/2022.
10. American Academy of Orthopedic Surgeons (AAOS). Compartment Syndrome. <https://orthoinfo.aaos.org/en/diseases--conditions/compartment-syndrome/>. Accessed 6/17/2022.
11. Undersea and Hyperbaric Medical Society. Compromised Skin Graft or Flaps. <https://www.uhms.org/12-compromised-grafts-and-flaps.html>. Accessed 6/27/2022.
12. Medline Plus. Crush Injury. <https://medlineplus.gov/ency/article/000024.htm>. Accessed 6/17/2022.
13. Medscape. Cyanide Toxicity. <https://emedicine.medscape.com/article/814287-overview>. Accessed 6/17/2022.
14. Harvard Health Publishing. Harvard Medical School: Decompression Sickness. Published: January, 2019. https://www.health.harvard.edu/a_to_z/decompression-sickness-a-to-z. Accessed 6/17/2022.
15. Shahab Qureshi M, FACP. Medscape: Clostridial Gas Gangrene. <https://emedicine.medscape.com/article/214992-overview>. Published 2017. Accessed 6/17/2022.
16. Medline Plus. Brain Abscess. <https://medlineplus.gov/ency/article/000783.htm>. Accessed 6/17/2022.
17. Medline Plus. Necrotizing Soft Tissue Infection. <https://medlineplus.gov/ency/article/001443.htm>. Accessed 6/17/2022.
18. American College of Hyperbaric Medicine. Radiation Necrosis. <http://achm.org/soft-tissue-radionecrosis/>. Accessed 6/17/2022.
19. Undersea and Hyperbaric Medical Society. Osteomyelitis (Refractory). <https://www.uhms.org/10-osteomyelitis-refractory.html>. Accessed 6/17/2022.
20. University of Maryland Medical Center. Anemia. <https://www.umms.org/ummc/health-services/shock-trauma/services/hyperbaric-medicine?si=ummc>. Accessed 6/17/2022.
21. Rauch SD. Clinical practice. Idiopathic sudden sensorineural hearing loss. *The New England journal of medicine*. 2008;359(8):833-840.
22. Hayes Inc. Topical Oxygen Therapy for Chronic Wound Healing. https://evidence.hayesinc.com/report/dir.topicaloxwound311?_ga=2.57821801.296794708.1590494915-1995923823.1570058383. Published 2017 (updated 2022). Accessed 6/17/2022.
23. Bennett MH, Lehm JP, Jepson N. Hyperbaric oxygen therapy for acute coronary syndrome. *The Cochrane database of systematic reviews*. 2015(7):Cd004818.
24. Health Evidence Review Commission. Health Evidence Review Commission (HERC) Coverage Guidance: Indications for Hyperbaric Oxygen Therapy. <https://www.oregon.gov/oha/HPA/DSI-HERC/EvidenceBasedReports/Indications-for-HBOT.pdf>. Published 2014. Accessed 6/17/2022.
25. U.S. Food and Drug Administration Consumer Alert: Hyperbaric Oxygen Therapy. <https://www.fda.gov/forconsumers/consumerupdates/ucm364687.htm>. Accessed 6/17/2022.

26. Bennett MH, Weibel S, Wasiak J, Schnabel A, French C, Kranke P. Hyperbaric oxygen therapy for acute ischaemic stroke. *The Cochrane database of systematic reviews*. 2014(11):Cd004954.
27. Eskes A, Vermeulen H, Lucas C, Ubbink DT. Hyperbaric oxygen therapy for treating acute surgical and traumatic wounds. *The Cochrane database of systematic reviews*. 2013(12):Cd008059.
28. Xiong T, Chen H, Luo R, Mu D. Hyperbaric oxygen therapy for people with autism spectrum disorder (ASD). *The Cochrane database of systematic reviews*. 2016;10:Cd010922.
29. National Institute for Health and Care Excellence. Autism: The Management and Support of Children and Young People on the Autism Spectrum. 2013.
<https://www.ncbi.nlm.nih.gov/pubmed/26065058>
30. National Collaborating Centre for Mental Health. National Institute for Health and Care Excellence: Guidance. In: *Autism: The Management and Support of Children and Young People on the Autism Spectrum*. Leicester (UK): British Psychological Society Copyright © National Collaborating Centre for Mental Health, 2013.; 2013.
31. Holland NJ, Bernstein JM, Hamilton JW. Hyperbaric oxygen therapy for Bell's palsy. *The Cochrane database of systematic reviews*. 2012(2):Cd007288.
32. Bennett MH, Trytko B, Jonker B. Hyperbaric oxygen therapy for the adjunctive treatment of traumatic brain injury. *The Cochrane database of systematic reviews*. 2012;12:Cd004609.
33. Bennett M, Best TM, Babul S, Taunton J, Lepawsky M. Hyperbaric oxygen therapy for delayed onset muscle soreness and closed soft tissue injury. *The Cochrane database of systematic reviews*. 2005(4):Cd004713.
34. Bennett MH, Stanford RE, Turner R. Hyperbaric oxygen therapy for promoting fracture healing and treating fracture non-union. *The Cochrane database of systematic reviews*. 2012;11:Cd004712.
35. Bennett MH, French C, Schnabel A, Wasiak J, Kranke P, Weibel S. Normobaric and hyperbaric oxygen therapy for the treatment and prevention of migraine and cluster headache. *The Cochrane database of systematic reviews*. 2015(12):Cd005219.
36. State of Colorado. Department of Labor and Employment. Division of Workers' Compensation. Rule17, Exhibit 6. Lower extremity injury medical treatment guidelines. Revised: January 19, 2016.
https://www.colorado.gov/pacific/sites/default/files/Rule_17_Exhibit_6_Lower_Extremity_Guideline.pdf. Accessed 4/30/2021.
37. Bennett M, Heard R. Hyperbaric oxygen therapy for multiple sclerosis. *CNS neuroscience & therapeutics*. 2010;16(2):115-124.
38. Phillips JS, Jones SE. Hyperbaric oxygen as an adjuvant treatment for malignant otitis externa. *The Cochrane database of systematic reviews*. 2013(5):Cd004617.
39. United States Department of Veterans Affairs. VA/DoD Clinical Practice Guidelines. Management of Posttraumatic Stress Disorder and Acute Stress Reaction 2017
<https://www.healthquality.va.gov/guidelines/MH/ptsd/>. Accessed 4/30/2021.
40. National Institute for Health and Clinical Excellence. The Prevention and Management of Pressure Ulcers in Primary and Secondary Care. 2014.
<https://www.ncbi.nlm.nih.gov/pubmed/25340232>
41. State of Colorado. Department of Labor and Employment. Division of Workers' Compensation. Shoulder injury medical treatment guidelines. Effective: November 2, 2008.
https://www.colorado.gov/pacific/sites/default/files/SI_Guideline%20with%20in-text%20references_0.pdf. Accessed 4/30/2021.

42. Bennett MH, Feldmeier J, Smee R, Milross C. Hyperbaric oxygenation for tumour sensitisation to radiotherapy. *The Cochrane database of systematic reviews*. 2012(4):Cd005007.
43. Xiao Y, Wang J, Jiang S, Luo H. Hyperbaric oxygen therapy for vascular dementia. *The Cochrane database of systematic reviews*. 2012(7):Cd009425.
44. Hayes PD, Alzuhir N, Curran G, Loftus IM. Topical oxygen therapy promotes the healing of chronic diabetic foot ulcers: a pilot study. *Journal of wound care*. 2017;26(11):652-660.
45. Agarwal V, Aroor S, Gupta N, Gupta A, Agarwal N, Kaur N. New Technique of Applying Topical Oxygen Therapy as a Cost-Effective Procedure. *The Indian journal of surgery*. 2015;77(Suppl 3):1456-1459.
46. Dissemond J, Kroger K, Storck M, Risse A, Engels P. Topical oxygen wound therapies for chronic wounds: a review. *Journal of wound care*. 2015;24(2):53-54, 56-60, 62-53.
47. Tawfick WA, Sultan S. Technical and clinical outcome of topical wound oxygen in comparison to conventional compression dressings in the management of refractory nonhealing venous ulcers. *Vascular and endovascular surgery*. 2013;47(1):30-37.
48. Belley R, Buteau D, Chateau-Degat ML. Efforts in trying to better define the level of evidence on topical pressurised oxygen therapy (TPOT) for wound healing. *International wound journal*. 2013;10(6):713-715.
49. Gordillo GM, Roy S, Khanna S, et al. Topical oxygen therapy induces vascular endothelial growth factor expression and improves closure of clinically presented chronic wounds. *Clinical and experimental pharmacology & physiology*. 2008;35(8):957-964.
50. Landau Z, Sommer A, Miller EB. Topical hyperbaric oxygen and low-energy laser for the treatment of chronic ulcers. *European journal of internal medicine*. 2006;17(4):272-275.
51. Kalliainen LK, Gordillo GM, Schlanger R, Sen CK. Topical oxygen as an adjunct to wound healing: a clinical case series. *Pathophysiology : the official journal of the International Society for Pathophysiology*. 2003;9(2):81-87.
52. Landau Z, Schattner A. Topical hyperbaric oxygen and low energy laser therapy for chronic diabetic foot ulcers resistant to conventional treatment. *The Yale journal of biology and medicine*. 2001;74(2):95-100.
53. Landau Z. Topical hyperbaric oxygen and low energy laser for the treatment of diabetic foot ulcers. *Archives of orthopaedic and trauma surgery*. 1998;117(3):156-158.
54. National Institute for Health and Care Excellence. Pressure ulcers: prevention and management. <https://www.nice.org.uk/guidance/cg179/resources/pressure-ulcers-prevention-and-management-pdf-35109760631749>. Published 2014. Accessed 5/13/2021.
55. U.S. Food and Drug Administration 510(k) Summary for Gulf Coast Hyperbarics, Inc. Rectangular Multiplace Hyperbaric Chamber (K082455). May 22, 2009. https://www.accessdata.fda.gov/cdrh_docs/pdf8/k082455.pdf. Accessed 6/17/2022.