


Policy and Procedure

PHARMACY PRIOR AUTHORIZATION POLICY AND CRITERIA ORPTCOTH030.1021	MISCELLANEOUS PRODUCTS INFUSION THERAPY SITE OF CARE
Effective Date: 1/1/2022 	Review/Revised Date: 11/19, 02/20, 05/20, 09/20, 08/21 (SNM/JL)
	P&T Committee Meeting Date: 10/19, 12/19, 04/20, 06/20, 08/20, 10/21
	Original Effective Date: 01/20
	Approved by: Oregon Region Pharmacy and Therapeutics Committee
Robert Gluckman, M.D. Chief Medical Officer	Page 1 of 6

SCOPE:

Providence Health Plan, Providence Health Assurance, Providence Plan Partners, and Ayn Health Solutions as applicable (referred to individually as “Company” and collectively as “Companies”).

APPLIES TO:

Commercial

Definitions:

1. **Site of Care** – the physical location where the infusion therapy is administered (e.g., an inpatient hospital, outpatient hospital-based infusion center, stand-alone infusion center, healthcare provider’s office, or home infusion)
2. **Alternative Site of Care** – any outpatient infusion site of care outside of an outpatient hospital-based infusion center (e.g., such as provider’s office or home infusion service providers)
3. **Approved Site of Care** - alternative sites of care or approved hospital-based infusion centers
4. **Unapproved Site of Care** – any site of care that has been deemed as medically unnecessary, including unapproved hospital based infusion centers that increase the cost of care compared to approved sites of care

POLICY CRITERIA:

COVERED USES:

The Company requires the infusion of certain medications (see [Table 1](#)) to be administered at an approved site of care, when an unapproved hospital-based infusion setting is determined to be no longer medically necessary.

REQUIRED MEDICAL INFORMATION:

1. Prior authorization for the medication must be obtained, if necessary. Refer to individual drug specific policies for clinical criteria.

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCOTH030**

**MISCELLANEOUS PRODUCTS
INFUSION THERAPY SITE OF CARE**

- a. For medications that require prior authorization for clinical criteria, the approval or denial of administration in an unapproved hospital outpatient setting is not indicative of approval or denial of the prior authorization for the medication based on clinical criteria.
2. The unapproved hospital-based outpatient infusion center may be considered medically necessary if one of the following criteria is met:
 - a. The patient has concomitant conditions or clinical history that may increase the risk of infusion reactions or drug specific adverse events, defined as one of the following:
 - i. Recent documented history of severe adverse drug reactions or anaphylaxis to prior treatments of the same or similar therapy.
 - ii. Concomitant complex medical conditions that may increase the risk of infusion reactions or complications to therapy. For example, the presence of antibodies that may increase the risk of infusion reactions, severely compromised cardiac and respiratory function.
 - iii. Use of multiple concurrent therapies of which one or more require infusion services at a higher level of care (e.g., cytotoxic chemotherapy, CAR-T given over same treatment period as requested medication)
 - iv. Chronic vascular access complications that require hospital-based interventions or equipment not available to home infusion providers
 - v. Mental health or cognitive changes that require increased level of care for the safe administration of infusions
 - b. The unapproved hospital based infusion center is deemed a more appropriate option, as defined by BOTH of the following criteria:
 - i. An approved site of care would require an additional 15 miles of travel from the member's home as compared to unapproved hospital based infusion center in the vicinity.
AND
 - ii. Home infusion services are not an option because the member's home is ineligible for infusion services. The eligibility of a member's home for home infusion can be affected by such factors as:
 1. The location of the member's home being outside of the infusion provider's service area, or
 2. Upon inspection, the home infusion provider considers the member's home to be unfit or unsafe for home infusion services.
3. The first 60 days after the drug authorization will be covered at an unapproved site of care, to accommodate for initial doses to be administered

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCOTH030**

**MISCELLANEOUS PRODUCTS
INFUSION THERAPY SITE OF CARE**

without delay to therapy. The purpose of the initial 60-day period is to allow for the determination of infusion tolerability at a higher level of care. This period will also allow for the timely submission and review of a prior authorization for the unapproved site of care, and the coordination of transition to an approved site, when the unapproved site of care has been determined to be not medically necessary.

EXCLUSION CRITERIA: N/A

AGE RESTRICTIONS:

This policy applies to those members who are 13 years of age and older.

PRESCRIBER RESTRICTIONS: N/A

COVERAGE DURATION:

Initial authorization and reauthorization will be approved for up to one year.

Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case

INTRODUCTION:

In the outpatient setting, infusion therapy was originally administered at outpatient hospital facilities, but improved technology now allows for safe and effective administration at alternative sites of care; such as medical clinics, stand-alone infusion providers, or home infusion. The practice of providing infusions outside of the hospital setting is well-established and accepted in clinical practice.

Alternative sites of care offer high-quality infusion services for patients and reduce the overall cost of care when compared to unapproved hospital outpatient infusion centers. When more than one medically appropriate site of care is available, the Companies will approve the use of the most affordable alternative.

Table 1. The administration of the following medications requires prior authorization for the use of an unapproved hospital-based infusion center when an approved site of care is an available treatment option.

HCPCS	Trade Name	Drug Name
<u>Inflammatory Conditions</u>		
J3262	Actemra	tocilizumab
J0490	Benlysta	belimumab

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCOTH030**

**MISCELLANEOUS PRODUCTS
INFUSION THERAPY SITE OF CARE**

J3380	Entyvio	vedolizumab
Q5103	Inflectra	infliximab-dyyb
J0129	Orencia	abatacept
J1745	Remicade	infliximab
Q5104	Renflexis	infliximab-abda
Q5121	Avsola	infliximab-axxq
J1602	Simponi Aria	golimumab
<u>Multiple Sclerosis</u>		
J2350	Ocrevus	ocrelizumab
<u>Miscellaneous Diseases</u>		
J1300	Soliris	eculizumab
J1303	Ultomiris	ravulizumab-cwvz
J0584	Crysvita	burosumab-twza
<u>Enzyme Replacement Therapy</u>		
J1458	Naglazyme	galsulfase
J1786	Cerezyme	imiglucerase
J3060	Elelyso	taliglucerase alfa
J3385	VPRIV	velaglucerase alfa
J2840	Kanuma	sebelipase alfa
<u>Intravenous Immune Globulin</u>		
J1566	Carimune NF, Gammagard S/D	
J1572	Flebogamma, Flebogamma DIF	
J1569	Gammagard	
J1568	Octagam	
J1555	Cuvitru	
J1459	Privigen	
J1557	Gammaplex	
J1561	Gammaked, Gamunex-C	
J1559	Hizentra	
J1558	Xembify	

POSITION STATEMENT:

In 2010, the National Home Infusion Association (NHIA) published a survey of home infusion providers that showed 829,000 patients were receiving therapy via home infusion. Of these patients, 1.24 million therapies were being administered via home

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCOTH030**

**MISCELLANEOUS PRODUCTS
INFUSION THERAPY SITE OF CARE**

infusion in 2008. Due to an aging population and an increase in chronic conditions requiring higher levels of treatment and monitoring, the U.S. home infusion market has continued to expand and provide clinical and safety data to support its use as a trusted site of care.

Patients receiving therapy via home infusion have been shown to have similar rates of positive clinical outcomes and adverse events as other sites of care. Patients have also reported preference rates of as high as 95% in favor of home infusion sites of care.

A retrospective chart review was done to analyze the incidence and management of infusion reactions to infliximab in an alternative care setting. A total of 796 patients with Crohn's disease or ulcerative colitis received a combined 5581 infliximab infusions with one home infusion provider between January 2014 and November 2016. Alternative care settings reviewed in the study were identified as either patient's home or the home infusion provider's infusion suite. Patients eligible for alternative care infusion were identified by their physician, and referred to either the home setting or an infusion suite. Use of premedication was determined by the referring physician, and all infusions were administered by a trained nurse following standardized protocols, who also did the post infusion monitoring for a minimum of 60min following the first 3 infusions and a minimum of 30min subsequently. In total, 109 infusion reactions (2% of all infusions) were recorded in 62 patients (7.8% of all patients). Of these reactions, 87 (79.8%) were acute, the majority of which were classified as mild (57.5%) or moderate in severity (31.0%). Ten infusions were associated with a severe reaction (11.5%, 0.2% of all infusions), and of these, 8 (9.2%, 0.1% of all infusions) resulted in an emergency room visit. The most common acute reaction was headache (23.0% of all acute IRs), followed by pruritus (14.9%); other common acute reactions included dyspnea (13.8%), flushing (13.8%), chest tightness/discomfort (11.5%), and nausea and/or vomiting (10.3%). In comparison, the REMICADE (infliximab) package insert reported infusion reactions in 18% of infusions, with serious infusion reactions occurring in <1% of patients.

All exceptions to the site of care medical necessity determination will be reviewed on a case-by-case basis.

The infused medication list included in this policy is subject to change.

REFERENCE/RESOURCES:

1. National Infusion Center Association. <https://infusioncenter.org> (Accessed 2019 Sept 2)

**PHARMACY PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCOTH030**

**MISCELLANEOUS PRODUCTS
INFUSION THERAPY SITE OF CARE**

2. Managed Healthcare Executive. Top ways to manage cost of infusible specialty therapies. <https://www.managedhealthcareexecutive.com/business-strategy/top-ways-manage-cost-infusible-specialty-therapies> (Accessed 2019 Sept 2)
3. National Home Infusion Association releases first-ever study of alternate-site infusion industry. www.nhia.org/press_release/pr101811.html (Accessed 2019 Sept 2)
4. Polinski JM, Kowal MK, Gagnon M et al. Home Infusion: Safe, clinically effective, patient preferred, and cost saving. *J of Del Sci Innov.* 2017;5:68-80.
5. Harris Williams & Co. Home infusion industry overview 2014. https://www.harriswilliams.com/sites/default/files/industry_reports/2014.6.12_home_infusion_industry_overview.pdf (Accessed 2019 Sept 2)
6. Remicade package insert. Horsham, PA: Janssen Biotech, Inc.; 2018 June.
7. Checkley LA, Kristofek L, Kile S et. Al. Incidence and management of infusion reactions to infliximab in an alternate care setting. *Dig Dis Sci.* 2019; 64(3):855-862.